

Strong convergence of Kleinian groups: the cracked eggshell

James W. Anderson

School of Mathematics, University of Southampton, Southampton SO17 1BJ England

Cyril Lecuire

Institut de Mathématiques de Toulouse, F-31062 Toulouse France

March 10, 2010

Abstract

In this paper we give a complete description of the set $\text{SH}(\pi_1(M))$ of discrete faithful representations of the fundamental group of a compact, orientable, hyperbolizable 3-manifold with incompressible boundary, equipped with the strong topology, with the description given in term of the end invariants of the quotient manifolds. As part of this description, we introduce coordinates on $\text{SH}(\pi_1(M))$ that extend the usual Ahlfors-Bers coordinates. We use these coordinates to show the local connectivity of $\text{SH}(\pi_1(M))$ and study the action of the modular group of M on $\text{SH}(\pi_1(M))$.

1 Introduction and statement of results

Kleinian groups have been studied since the late 19th century in the work of Poincaré and Fricke and Klein, and more extensively since the work of Ahlfors and Bers in the 1960s and Thurston in the 1970s and 1980s. In this paper, we consider a particular aspect of the basic question of understanding the behavior of sequences of Kleinian groups.

There are two standard notions of convergence for a sequence of Kleinian groups. The first is algebraic convergence, which is convergence on generators. The second is geometric convergence, which is convergence of the quotient hyperbolic 3-manifolds. The topology of algebraic convergence of the set of isomorphic, finitely generated Kleinian groups is well understood. In particular, there is a correspondence between connected components of this set of Kleinian groups and marked homotopy classes of compact 3-manifolds, up to a natural equivalence. Geometric convergence is considerably less well-behaved, as isomorphic, finitely generated Kleinian groups may converge geometrically to an infinitely generated Kleinian group.

Strong convergence of Kleinian groups combines these two different notions, so that a sequence of Kleinian groups converges strongly to a Kleinian group if it converges to that Kleinian group both algebraically and geometrically. Thurston proposed a picture of what the space of Kleinian groups with the strong topology looks like, which we consider here.

Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $D(\pi_1(M))$ denote the set of discrete faithful representations $\rho : \pi_1(M) \rightarrow \text{PSL}(2, \mathbb{C})$

equipped with the topology of algebraic convergence (formal definitions are given in Section 2). The group $\text{Isom}^+(\mathbb{H}^3) \simeq \text{PSL}(2, \mathbb{C})$ of orientation-preserving isometries of hyperbolic 3-space acts on $D(\pi_1(M))$ by conjugation. We denote the quotient by $\text{AH}(\pi_1(M))$. Note that a sequence $\{[\rho_n]\} \subset \text{AH}(\pi_1(M))$ converges to $[\rho] \in \text{AH}(\pi_1(M))$ if there is a sequence $\{h_n\} \subset \text{PSL}(2, \mathbb{C})$ so that $\{h_n \rho_n h_n^{-1}\}$ converges to ρ in $D(\pi_1(M))$. When there is no possibility of confusion, we refer to elements of both $D(\pi_1(M))$ and $\text{AH}(\pi_1(M))$ as representations, even though the elements of the latter set are formally equivalence classes of representations.

By considering the quotient manifold $M_\rho = \mathbb{H}^3/\rho(\pi_1(M))$ corresponding to a representation ρ , we can identify $\text{AH}(\pi_1(M))$ with the set of hyperbolic 3-manifolds homotopy equivalent to M up to isometry, where the homotopy equivalence induces the given representation. A sequence $\{\rho_n\}$ in $\text{AH}(\pi_1(M))$ converges strongly to a representation ρ_∞ if $\{\rho_n\}$ converges to ρ_∞ algebraically and if $\{\rho_n(\pi_1(M))\}$ converges geometrically to $\rho_\infty(\pi_1(M))$. We denote by $\text{SH}(\pi_1(M))$ the set $\text{AH}(\pi_1(M))$ equipped with the strong topology. Though we do not use this description, one can identify $\text{SH}(\pi_1(M))$ with the space of hyperbolic manifolds homotopy equivalent to M (up to isometry) equipped with a marked pointed Hausdorff-Gromov topology.

By Bonahon, see [Bon], each representation $\rho \in \text{AH}(\pi_1(M))$ is tame, namely the corresponding quotient manifold $M_\rho = \mathbb{H}^3/\rho(\pi_1(M))$ is homeomorphic to the interior of a compact 3-manifold. Combining this result with the uniqueness of the compact core, see [MMS], it follows that the topology of the quotient manifold does not change under strong convergence (compare with [CaM]). Namely if $\{\rho_n\}$ converges strongly to ρ_∞ , then $M_\infty = \mathbb{H}^3/\rho_\infty(\pi_1(M))$ is homeomorphic to $M_n = \mathbb{H}^3/\rho_n(\pi_1(M))$ for n sufficiently large. Using the Ending Lamination Theorem, see [Mi2] and [BCM], and Thurston's Double Limit Theorem, see [Th3], one can then see that the connected components of $\text{SH}(\pi_1(M))$ are in 1-to-1 correspondence with the set of marked homeomorphism types of 3-manifolds homotopy equivalent to M (we see that this fact can be deduced from our results as well). In particular, as a first point of contrast with $\text{AH}(\pi_1(M))$, we show that $\text{SH}(\pi_1(M))$ does not have the bumping phenomenon described for $\text{AH}(\pi_1(M))$ for many M in [AnC].

In addition to the existence of bumping, another disturbing property of $\text{AH}(\pi_1(M))$ is that it may not even be locally connected, see [Br]. We will show that this does not happen with the strong topology.

Theorem A. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Then the space $\text{SH}(\pi_1(M))$ is locally connected.*

Even when M is such that $\text{AH}(\pi_1(M))$ is locally connected, the space $\text{AH}(\pi_1(M))$ may have so called self-bumping points, see [McM] and [BrH]. We will see that this kind of phenomenon does not appear in $\text{SH}(\pi_1(M))$.

Theorem B. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\rho \in \text{SH}(\pi_1(M))$ be a representation uniformizing M . Then any neighborhood of ρ contains a neighborhood $\mathcal{V} \subset \text{SH}(\pi_1(M))$ of ρ such that $\mathcal{V} \cap \text{int}(\text{SH}(\pi_1(M)))$ is connected.*

The reason why self-bumping points may be present in $\text{AH}(\pi_1(M))$ and not in $\text{SH}(\pi_1(M))$ is that some sequences of representations that converge in $\text{AH}(\pi_1(M))$ do not converge in $\text{SH}(\pi_1(M))$. This has the following consequence:

Lemma 1.1. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Then the space $\text{SH}(\pi(M))$ is not locally compact.*

We denote by $\text{AH}(M) \subset \text{AH}(\pi_1(M))$ the set of representations ρ whose quotient manifold $\mathbb{H}^3/\rho(\pi_1(M))$ is homeomorphic to the interior of M by a homeomorphism that induces ρ . This set is contained in the closure of the component of the interior of $\text{AH}(\pi_1(M))$ containing the minimally parabolic Kleinian groups Γ for which M_Γ is homeomorphic to the interior of M (this can be deduced from [Mar], [Su], [BCM]). Let $\text{SH}(M)$ denote the set $\text{AH}(M)$ with the strong topology. Let $\text{Mod}(M)$ be the group of isotopy classes of orientation-preserving diffeomorphisms of M . When $M = S \times I$ is a trivial I -bundle, Kerckhoff and Thurston, see [KeT], have shown that the action of $\text{Mod}(M) = \text{MCG}(S)$ on $\text{AH}(M)$ is not properly discontinuous. On $\text{SH}(M)$ we have the following result.

Theorem C. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Assume that M is not an I -bundle over a closed surface. Then the action of $\text{Mod}(M)$ on $\text{SH}(M)$ is properly discontinuous.*

Furthermore, we show that when M is a trivial I -bundle, this action has fixed points.

1.1 Outline of the paper

Thurston [Th2] gives the following conjectural description of $\text{AH}(M)$ and $\text{SH}(M)$. When M does not contain an essential annulus, $\text{AH}(M)$ is homeomorphic to a closed ball, like an hard-boiled egg with its shell, whereas $\text{SH}(M)$ is obtained by thoroughly cracking the eggshell on a hard surface. In order to get a precise description of this cracked eggshell, we introduce some coordinates for $\text{SH}(M)$.

Say that a representation $\rho \in \text{AH}(M)$ *uniformizes* M . By a result of Scott, see [Sc], M_ρ has a compact core K homeomorphic to M . Choosing K to intersect the cusps of M_ρ in annuli or tori, see [McC], we may define the ends of M_ρ to be the complementary regions of $K \cup \text{cusps}$. Such an end is *geometrically infinite* if it can be chosen to be contained in the convex core C_ρ of M_ρ and *geometrically finite* otherwise. We define the *conformal end invariants* of ρ to consist of the hyperbolic metrics associated to the conformal structure at infinity of each geometrically finite end and of the ending laminations associated to its geometrically infinite ones. Such an invariant (F, m, L) consists of a complete hyperbolic metric m on an open subsurface F of ∂M and of a geodesic lamination L that "almost fills" its complement; we call it a *gallimaufry* on M or on ∂M (see definition in section 2.8). We put on the set of gallimaufries a topology that extends the usual topology of the Teichmüller space. We say that the gallimaufry (F, m, L) is *doubly incompressible* if there is a transverse measure λ supported by L and $\eta > 0$ such that for any essential annulus or Möbius band $(E, \partial E) \subset (M, \partial M - F)$ we have $i(\partial E, \lambda) \geq \eta$. We prove the following result:

Theorem D. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. The ending map that associates to a representation uniformising M its end invariants is a homeomorphism from $\text{SH}(M)$ into the set of doubly incompressible gallimaufries on ∂M .*

By the proof of the Ending Lamination Theorem, see [Mi2] and [BCM], the map associating its ending gallimaufry to a (conjugacy class of) representation is 1-to-1. Thus we only need to show that the ending map defined in Theorem D is proper and continuous.

We then prove that the set of doubly incompressible gallimaufries equipped with its topology is locally connected, does not have self-bumping points, and is not locally compact.

In section 2, we give the definitions and some basic results that are used in the paper. In particular, we define the space of gallimaufries and describe some of its properties. The proof that the ending map is proper is divided in two. In section 3 we show that given a sequence of representations whose end invariants converge to a doubly incompressible gallimaufry, a sub-sequence converges algebraically. This can be viewed as a refinement of Thurston's Double Limit Theorem. The proof mixes arguments of Thurston's original proof, see [Th3], arguments of Otal's proof, see [Ot2], and a cut and paste operation. In section 4 we show that the algebraically convergent sub-sequence provided in the preceding section actually converges strongly, by studying the behaviour of the convex cores of the sequence and concluding that the limit sets converge in the Hausdorff topology. In section 5 we show that the ending map is continuous. The main difficulty here is in handling the geometrically infinite ends of the limit. This will be dealt with by proving a relative version of Bonahon's Intersection Lemma, see [Bon]. At this point we have proved Theorem D and Lemma 1.1. In section 6, we prove Theorems A and B by constructing appropriate paths in the space of doubly incompressible gallimaufries. In section 7 we prove Theorem C.

2 Background material and definitions

The purpose of this Section is to provide the background material we use in this paper. We often pass to sub-sequences; unless otherwise stated, we will without further comment use the same notation for the sub-sequence as for the original sequence.

2.1 Kleinian groups and 3-manifolds

Standard sources for material on Kleinian groups and hyperbolic 3-manifolds are [Th1], [MaT] and [Ka]. A *Kleinian group* is a discrete subgroup of the group $\mathrm{PSL}(2, \mathbb{C})$ of all orientation-preserving isometries of hyperbolic 3-space $\mathbb{H}^3$. Throughout this paper, we assume that Kleinian groups are torsion-free and non-elementary, so that they contain a non-abelian free subgroup.

A *hyperbolizable 3-manifold* is an orientable 3-manifold that admits a complete Riemannian metric all of whose sectional curvatures are equal to -1 . A 3-manifold endowed with such a metric is a *hyperbolic 3-manifold*. It follows that an orientable hyperbolic 3-manifold N can be expressed as the quotient $N = \mathbb{H}^3/\Gamma$ for a Kleinian group Γ , and that Γ is unique up to conjugacy in $\mathrm{PSL}(2, \mathbb{C})$.

An (orientation-preserving) isometry of $\mathbb{H}^3$ extends to a conformal transformation on the boundary at infinity $\partial_\infty \mathbb{H}^3 = \widehat{\mathbb{C}}$ of $\mathbb{H}^3$. The *domain of discontinuity* Ω_Γ of a Kleinian group Γ is the largest open subset of $\widehat{\mathbb{C}}$ on which the action of Γ is properly discontinuous.

The *limit set* $\Lambda_\Gamma \subset \widehat{\mathbb{C}}$ of Γ is the complement of Ω_Γ . Let $H_\Gamma \subset \mathbb{H}^3$ be the *convex hull* in $\mathbb{H}^3$ of Λ_Γ , which is the smallest non-empty convex subset of $\mathbb{H}^3$ invariant under the action of Γ . The *convex core* C_Γ of Γ is the quotient of H_Γ by the action of Γ .

A Kleinian group Γ is *geometrically finite* if the unit neighbourhood of its convex core C_Γ has finite volume in $\mathbb{H}^3/\Gamma$. A Kleinian group Γ is *convex co-compact* if its convex core C_Γ is compact, or equivalently, if it is geometrically finite and contains no parabolic elements. A Kleinian group is *minimally parabolic* if every parabolic isometry belongs to a rank two free abelian subgroup.

A compact 3-manifold is *hyperbolizable* if there exists a Kleinian group Γ so that the interior $\text{int}(M)$ of M is homeomorphic to the quotient manifold $\mathbb{H}^3/\Gamma$. Note that, under this definition, a hyperbolizable 3-manifold M is necessarily *irreducible* (so that every embedded 2-sphere in M bounds a 3-ball in M), orientable, and *atoroidal* (so that every incompressible torus T in M is homotopic into ∂M).

An embedded compact sub-manifold W in a 3-manifold M is *incompressible* if $\pi_1(W)$ is infinite and if the inclusion $W \hookrightarrow M$ induces an injective map on fundamental groups. A compact 3-manifold has *incompressible boundary* if each component of ∂M is incompressible.

For a compact 3-manifold M , let $\partial_{\chi < 0} M$ denote the union of the components of ∂M of negative Euler characteristic, so that $\partial_{\chi < 0} M$ consists of ∂M with any boundary tori removed.

Let $F \subset \partial M$ be an incompressible compact surface. An *essential annulus* (or Möbius band) E in (M, F) is a properly embedded incompressible annulus (or Möbius band) $(E, \partial E) \subset (M, F)$ that cannot be homotoped into F by a homotopy fixing ∂E . An *essential disc* D is a properly embedded disc $(D, \partial D) \subset (M, F)$ that cannot be homotoped into F by a homotopy fixing ∂D .

2.2 Discrete faithful representations

Let M be a compact, orientable, hyperbolizable 3-manifold. A *discrete faithful representation* of $\pi_1(M)$ into $\text{PSL}(2, \mathbb{C})$ is an injective homomorphism $\rho : \pi_1(M) \rightarrow \text{PSL}(2, \mathbb{C})$ whose image $\rho(\pi_1(M))$ is a Kleinian group. A discrete faithful representation ρ of $\pi_1(M)$ is *geometrically finite* if the image group $\rho(\pi_1(M))$ is geometrically finite, and is *convex co-compact* if $\rho(\pi_1(M))$ is convex co-compact.

Let $\rho : \pi_1(M) \rightarrow \text{PSL}(2, \mathbb{C})$ be a discrete faithful representation. Assume moreover that the quotient manifold $M_\rho = \mathbb{H}^3/\rho(\pi_1(M))$ is homeomorphic to $\text{int}(M)$ by a homeomorphism $f : \text{int}(M) \rightarrow M_\rho = \mathbb{H}^3/\rho(\pi_1(M))$ that induces ρ , so that $\rho = f_*$. Under these assumptions, say that ρ *uniformises* M .

Choose now a pairwise disjoint set of horoballs in $\mathbb{H}^3$ so that each horoball is centered at a parabolic fixed point of $\rho(\pi_1(M))$, and there is a horoball centered at the fixed point of each parabolic subgroup of $\rho(\pi_1(M))$ and hence invariant under the corresponding parabolic subgroup. It is a standard application of the Margulis lemma that such a set of horoballs exists. The *cuspidal part* of M_ρ is the quotient of such a collection of horoballs. Where relevant, we will assume that we have made a convenient choice of such a collection of horoballs.

Let $D(\pi_1(M))$ be the space of all discrete faithful representations of $\pi_1(M)$ into $\text{PSL}(2, \mathbb{C})$. By choosing a fixed set of p generators $g_1, \dots, g_p$ for $\pi_1(M)$, we can realize $D(\pi_1(M))$ as a

subspace of $(\mathrm{PSL}(2, \mathbb{C}))^p$ by the identification $\rho \rightarrow (\rho(g_1), \dots, \rho(g_p))$; the topology thus obtained is called the *algebraic topology* or the *topology of algebraic convergence*. By [Jor], $D(\pi_1(M))$ is closed in this topology. In this paper, we do not work with all of $D(\pi_1(M))$, but rather with the connected components of its interior; for a full description of $D(\pi_1(M))$ in light of the proof of the Ending Lamination Theorem, see [ACC]. The representations contained in a given connected component of the interior of $D(\pi_1(M))$ are the geometrically finite minimally parabolic representations uniformising a given compact, orientable, hyperbolizable 3-manifold M' homotopy equivalent to M , by work of Ahlfors, Bers, Kra, Marden, Maskit, Sullivan, and Thurston. Such a component of $\mathrm{int}(D(\pi_1(M)))$ is uniquely defined by the given manifold M' .

If we instead consider the Chabauty topology on subgroups of $\mathrm{PSL}(2, \mathbb{C})$, we get the *geometric topology* or the *topology of geometric convergence*. A sequence $\{\rho_n\} \subset D(\pi_1(M))$ converges geometrically to a Kleinian group G_∞ if every accumulation point a of every sequence $\{a_n \in \rho_n(\pi_1(M))\}$ lies in G_∞ for and if every element a of G_∞ is the limit of a sequence $\{a_n \in \rho_n(\pi_1(M))\}$. Note that the limit manifold $\mathbb{H}^3/G_\infty$ is not necessarily homeomorphic to $\mathrm{int}(M)$, and in fact the geometric limit G_∞ of a sequence of (isomorphic) finitely generated Kleinian groups can be infinitely generated.

The sequence $\{\rho_n\} \subset D(\pi_1(M))$ converges strongly if $\{\rho_n\}$ converges algebraically to some representation ρ_∞ and if $\{\rho_n(\pi_1(M))\}$ converges geometrically to $\rho_\infty(\pi_1(M))$. As we commented on in the introduction, strong convergence preserves the topology of the quotient manifold.

Lemma 2.1. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\{\rho_n\}$ be a sequence of representations uniformising M that converges strongly to a representation ρ_∞ . Then ρ_∞ uniformises M .*

Proof. By the proof of Marden's tameness conjecture, [Ag] or [CaG], see also [Bon], there is a compact 3-manifold M' such that $M_\infty = \mathbb{H}^3/\rho_\infty(\pi_1(M))$ is homeomorphic to $\mathrm{int}(M')$. Consider a compact core K for M_∞ , which we can choose to be homeomorphic to M' . Since $\{\rho_n(\pi_1(M))\}$ converges geometrically to $\rho_\infty(\pi_1(M))$, there are points x_n , $n \in \mathbb{N} \cup \infty$, and maps $\phi_n : M_\infty \rightarrow M_n$ with $\phi_n(x_\infty) = x_n$ such that the restriction of ϕ_n to $B(x_\infty, R_n)$ is q_n -bilipschitz with $R_n \rightarrow \infty$ and $q_n \rightarrow 1$, see [MaT]. For n sufficiently large, we have $K \subset B(x_\infty, R_n)$. By construction $\phi_n(K)$ is a compact core for M_n . By [MMS], any compact core for M_n is homeomorphic to K . It follows that M_n is homeomorphic to M_∞ for n sufficiently large. $\square$

On $\mathrm{int}(D(\pi_1(M)))$, the algebraic and strong topologies are equivalent.

The group $\mathrm{PSL}(2, \mathbb{C})$ acts on $D(\pi_1(M))$ by conjugation. We denote by $\mathrm{AH}(\pi_1(M))$ the quotient of $D(\pi_1(M))$ by $\mathrm{PSL}(2, \mathbb{C})$ endowed with the algebraic topology. We denote by $\mathrm{SH}(\pi_1(M))$ the quotient of $D(\pi_1(M))$ by $\mathrm{PSL}(2, \mathbb{C})$ endowed with the strong topology. The interiors of both $\mathrm{AH}(\pi_1(M))$ and $\mathrm{SH}(\pi_1(M))$, with their respective topologies, are the quotient of $\mathrm{int}(D(\pi_1(M)))$ by $\mathrm{PSL}(2, \mathbb{C})$.

Let $D(M)$ be the set of representations $\rho \in D(\pi_1(M))$ that uniformise M . Recall that this means that the quotient manifold $M_\rho = \mathbb{H}^3/\rho(\pi_1(M))$ is homeomorphic to $\mathrm{int}(M)$

by a homeomorphism $f : \text{int}(M) \rightarrow M_\rho = \mathbb{H}^3/\rho(\pi_1(M))$ that induces ρ . (Hence, the interior of $D(M)$ consists of quasiconformal deformations of a given minimally parabolic Kleinian group uniformizing M .) We denote by $\text{AH}(M)$, respectively $\text{SH}(M)$, the quotient of $D(\pi_1(M))$ by $\text{PSL}(2, \mathbb{C})$ endowed with the algebraic topology, respectively the strong topology. When ∂M is incompressible, it follows from the work of Ahlfors and Bers that $\text{int}(\text{AH}(M)) = \text{int}(\text{SH}(M))$ is homeomorphic to the Teichmüller space $\mathcal{T}(\partial_{\chi < 0} M)$; in particular it is a topological ball.

It follows from Lemma 2.1 that for any topological manifold M , $\text{SH}(M)$ is a closed set and is disjoint from any other set $\text{SH}(M') \subset \text{SH}(\pi_1(M))$. Hence to understand $\text{SH}(\pi_1(M))$, we only need to understand each of the $\text{SH}(M')$ separately as M' varies over all the 3-manifolds homotopy equivalent to M .

An $\mathbb{R}$ -tree is a path metric space such that any two points can be joined by a unique geodesic arc. Let $\mathcal{T}$ be an $\mathbb{R}$ -tree and let G be a group acting on $\mathcal{T}$. We say that the action is *small* if the edge stabilizers are virtually cyclic. We say that the action is *minimal* if no proper subtree of $\mathcal{T}$ is invariant under the action of G . We will normally consider actions of $\mathbb{R}$ -trees that are both small and minimal.

Morgan and Shalen [MoS1] introduced a compactification of $\text{AH}(M)$ by small minimal isometric actions of $\pi_1(M)$ on $\mathbb{R}$ -trees. Consider an element g of $\pi_1(M)$. For $\rho \in \text{AH}(M)$ we denote by $\ell_\rho(g)$ the translation distance of $\rho(g)$ (which is 0 if the isometry is parabolic). Given an isometric action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$, we denote by $\ell_{\mathcal{T}}(g)$ the translation distance of the action of g on $\mathcal{T}$. A sequence of representations $\{\rho_n\} \subset \text{AH}(M)$ tends to a minimal isometric action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$ if there are $\varepsilon_n \rightarrow 0$ such that $\varepsilon_n \ell_{\rho_n}(g) \rightarrow \ell_{\mathcal{T}}(g)$ for any element g of $\pi_1(M)$, see [Ot1]. Multiplying the sequence ε_n by a given constant gives us an isometric action on another $\mathbb{R}$ -tree which is homothetic to the first one. Hence $\text{AH}(M)$ has thus been compactified by actions on $\mathbb{R}$ -trees up to homothety.

Replacing $\mathbb{H}^3$ by $\mathbb{H}^2$ we get a compactification of the Teichmüller space which by Skora's Theorem, see [Sk], is equivalent to Thurston's compactification by projective measured geodesic laminations (see section 2.6).

2.3 Geodesic laminations

Standard sources for material on geodesic laminations are [Pe] and [Ot2, Appendice]. Note that unless otherwise explicitly stated, a surface of negative Euler characteristic will be equipped with a complete hyperbolic metric of finite area.

A *geodesic lamination* L on a closed hyperbolic surface S is a compact set that is the (non-empty) disjoint union of complete embedded geodesics. Note that this definition can be made independent of the choice of metric on S , see [Ot2, Appendice] for example. A geodesic lamination is *minimal* if it does not contain a geodesic lamination as a proper subset. A minimal lamination is either a simple closed geodesic or an *irrational lamination*. A leaf of a geodesic lamination is *recurrent* if it lies in a minimal sublamination. A geodesic lamination is the disjoint union of finitely many minimal sublaminations and finitely many non-recurrent leaves. The *recurrent part* of a geodesic lamination is the union of its recurrent leaves. It is itself a geodesic lamination. A *multi-curve* is a union of disjoint simple closed geodesics.

We say that two geodesic laminations L and L' *cross* if at least one leaf of L transversely intersects a leaf of L' .

Let F be a compact surface with boundary. We define a geodesic lamination in F similarly as in the case of a closed surface by considering a hyperbolic metric with geodesic boundary on F (again the definition can be made independent of the metric). A geodesic lamination $L \subset F$ is *peripheral* if L is a simple closed curve freely homotopic to a component of ∂F .

Let $L \subset S$ be a connected geodesic lamination that is not a simple closed curve. The *warren* $W(L)$ of L is the smallest subsurface of S with geodesic boundary containing L . Notice that we may have $W(L) = S$. It is not hard to see that $W(L)$ contains finitely many simple closed curves that are disjoint from L , see [Le1, §2.4]. Removing from $W(L)$ an annulus around each such curve, we get the *surface embraced by L* that we denote by $S(L)$. When L is a simple closed curve $S(L)$ is an (open) annular neighbourhood of L . In the particular case that L is a non-connected geodesic lamination then $S(L)$ is the disjoint union of the $S(L_i)$ as L_i runs through the connected components of L . Notice that $S(L)$ is an open surface (i.e. without boundary).

A *measured geodesic lamination* λ consists of a geodesic lamination $|\lambda|$ and a transverse measure on $|\lambda|$. Any arc $k \cong [0, 1]$ embedded in S transverse to $|\lambda|$, such that $\partial k \subset S - |\lambda|$, is endowed with a transverse measure $d\lambda$ such that:

- the support of $d\lambda|_k$ is $|\lambda| \cap k$;
- if an arc k' can be homotoped to k by a homotopy respecting $|\lambda|$ then $\int_k d\lambda = \int_{k'} d\lambda$.

If λ is a measured geodesic lamination, then its support $|\lambda|$ contains only recurrent leaves. Two measured geodesic laminations cross if their supports cross.

We denote by $\mathcal{ML}(S)$ the space of measured geodesic laminations on S endowed with the weak* topology on transverse measures. If γ is a weighted simple closed curve with weight $w(\gamma)$ and λ is a measured geodesic lamination transverse to γ , the *intersection number* $i(\lambda, \gamma)$ is defined by $i(\lambda, \gamma) = w(\gamma) \int_{|\gamma|} d\lambda$. Weighted simple closed curves are dense in $\mathcal{ML}(S)$ and so i extends to a continuous function $i : \mathcal{ML}(S) \times \mathcal{ML}(S) \rightarrow \mathbb{R}$, see [Bon].

Given a complete hyperbolic metric s on S , the length of a weighted simple closed curve with support c and weight $w \in \mathbb{R}$ is $w\ell_s(c)$, where $\ell_s(c)$ is the length of c with respect to s . This length function extends continuously to a function $\ell_s : \mathcal{ML}(S) \rightarrow \mathbb{R}$ called the *length function*, see [Bon].

This definition can be extended to define the length of a measured geodesic lamination in a hyperbolic 3-manifold as follows. Let M be a hyperbolic 3-manifold with boundary. We are not interested in curves lying in a torus component of ∂M , so we use the notation $\mathcal{ML}(\partial M)$ for $\mathcal{ML}(\partial_{\chi < 0} M)$. Let S be compact subsurface of $\partial_{\chi < 0} M$. Let $\rho \in \text{AH}(M)$ be a representation uniformizing M and let $c \subset S$ be a simple closed curve. Denote by c^* the closed geodesic in $M_\rho = \mathbb{H}^3 / \rho(\pi_1(S))$ in the free homotopy class defined by c , if such a geodesic exists. We denote by $\ell_\rho(c^*)$ the length of c^* with respect to the hyperbolic metric on M_ρ . When $\rho(c)$ is

a parabolic isometry, we take $\ell_\rho(c^*) = 0$. This allows us to define the length of a weighted multi-curve. Using the density of weighted multi-curves in $\mathcal{ML}(\partial M)$, we can then define the length $\ell_\rho(\lambda^*)$ of a measured geodesic lamination $\lambda \in \mathcal{ML}(\partial M)$.

We can associate to a measured geodesic lamination β on a hyperbolic surface S an action of $\pi_1(S)$ on an $\mathbb{R}$ -tree $\mathcal{T}_\beta$ dual to β , see [Ot2]. If c is a simple closed curve, we denote by $\ell_\beta(c)$ the translation distance of an isometry of $\mathcal{T}_\beta$ corresponding to c . The action of $\pi_1(S)$ on $\mathcal{T}_\beta$ we get in this way satisfies $\ell_\beta(c) = i(\beta, c)$. Notice that this property completely defines the action of $\pi_1(S)$ on $\mathcal{T}_\beta$, see [Ot2].

Let S be a connected hyperbolic surface and let $q : \mathbb{H}^2 \rightarrow S$ be the covering projection. Let $L \subset S$ be a geodesic lamination and let $\pi_1(S) \curvearrowright \mathcal{T}$ be a minimal action of $\pi_1(S)$ on an $\mathbb{R}$ -tree $\mathcal{T}$. Then L is *realized* in $\mathcal{T}$ if there is a continuous equivariant map $\mathbb{H}^2 \rightarrow \mathcal{T}$ whose restriction to any lift of a leaf of L is injective. When S is a component of the boundary of 3-manifold M , we extend this definition to actions of $\pi_1(M)$ on $\mathbb{R}$ -trees in the following way. Given an action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$, we use the map $i_* : \pi_1(\partial M) \rightarrow \pi_1(M)$ induced by the inclusion to get an action of $\pi_1(\partial M)$ on $\mathcal{T}$ (which is still small when ∂M is incompressible). By saying that L is realized in $\mathcal{T}$, we actually mean that L is realized in the minimal tree invariant under the action of $i_*(\pi_1(S))$ on $\mathcal{T}$, equipped with the action of $\pi_1(S)$.

A simple closed curve c in a compact (i.e. closed or compact with boundary) surface S is *essential* if c does not bound a disc in S . A compact subsurface $F \subset S$ is *essential* if any simple closed curve $c \subset F$ that bounds a disc in S bounds a disc in F .

2.4 Train tracks

Consider a compact surface S endowed with a complete hyperbolic metric of finite area. The purpose of this section is to introduce the notion of a *train track*, which is an object on a surface used to combinatorially encode measured geodesic laminations. Specifically, a *train track* τ in S is the union of finitely many "rectangles" b_i , called the *branches*. In a rectangle $[0, 1] \times [0, 1]$ we call a segment $\{p\} \times [0, 1] \subset [0, 1] \times [0, 1]$ a *vertical* segment and a segment $[0, 1] \times \{p\} \subset [0, 1] \times [0, 1]$ a *horizontal* segment. The 2 extremal vertical segments are the *vertical sides*. The branches of a train track satisfy:

- a branch b_i is the image of a rectangle $[0, 1] \times [0, 1]$ under a smooth map whose restriction to $]0, 1[\times]0, 1[$ is an embedding;
- the union of the double points of a branch is either empty or a non-degenerate vertical segment;
- given a pair of branches and the corresponding rectangles and maps, the intersection of the images is either empty or a non-degenerate segment;
- given the collection of rectangles and smooth maps producing the branches, any connected component of the union of the images of the vertical sides is a simple arc embedded in S .

The images of the vertical segments $\{p\} \times [0, 1]$ are the *ties*. A maximal connected union of ties lying in different branches is a *switch*. A sub-track of a train track τ is a train track all of whose branches are branches of τ .

A geodesic lamination L is *carried by a train track* τ when there is a hyperbolic metric m on S such that the m -geodesic lamination L lies in τ and is transverse to the ties. A geodesic lamination L is *minimally carried* by τ if no proper sub-track of τ carries L . A measured geodesic lamination λ is *carried* by a train track τ if its support $|\lambda|$ is carried by τ . Associating to each branch b of τ the transverse measure $\lambda(b)$ of a tie, we get a *weight system* for τ . A weight system is a function $\{\text{branches}\} \rightarrow \mathbb{R}^+$ satisfying the switch conditions: each switch is 2-sided and the sum of the weights of the branches on one side is equal to the sum on the other side. If a train track τ minimally carries a measured geodesic lamination, then there is a bijection between the set of measured geodesic laminations carried by τ and the weight systems for τ , see [Pe].

2.5 End invariants

Consider a discrete faithful representation $\rho : \pi_1(M) \rightarrow \text{PSL}(2, \mathbb{C})$ that uniformizes M . By a result of Scott, the quotient manifold M_ρ has a compact core K homeomorphic to M . We can choose K to intersect a chosen collection $\mathcal{C}$ of cusps of M_ρ (see Section 2.2) in annuli or tori. The *ends* of M_ρ are (equivalence classes of) the complementary regions in M_ρ of $K \cup \mathcal{C}$. To each end we associate an open surface $F \subset \partial K \approx \partial M$, and the end is homeomorphic to $F \times [0, \infty)$, by the Tameness Theorem.

Denote the convex core of M_ρ by C_ρ . An end of M_ρ , as defined above, is *geometrically infinite* if it is contained in the convex core C_ρ of M_ρ (up to a compact piece) and *geometrically finite* otherwise. We note that a Kleinian group is geometrically finite by the definition given earlier in Section 2.1 if and only if it has finitely many ends, each of which is geometrically finite by this definition.

The quotient $\Omega_\rho / \rho(\pi_1(M))$ of the domain of discontinuity is a Riemann surface of finite type and adds a natural conformal boundary to the open manifold M_ρ . This yields a natural embedding $\Omega_\rho / \rho(\pi_1(M)) \hookrightarrow \partial M$ well defined up to homotopy. To each geometrically finite end $E = F \times [0, \infty)$ is associated the component of $\Omega_\rho / \rho(\pi_1(M))$ homeomorphic to F . Thus we get a point in the Teichmüller space $\mathcal{T}(F)$; this is the end invariant associated to this end.

Let $E \approx F \times [0, \infty)$ be a geometrically infinite end of M_ρ . Thurston associated to E , see [Th1], a minimal geodesic lamination L on F defined as follows. Consider a sequence $\{c_n\} \subset F$ of simple closed curves whose geodesic representatives $c_n^* \subset E \subset M_\rho$ exit every compact subset of E . Make c_n into a geodesic lamination by using the counting measure. The existence of such a sequence of curves is a non-trivial fact proved in [Bon]. Extract a subsequence such that $\{c_n\}$ converges in $\mathcal{PML}(F)$ to a projective measured geodesic lamination $[\lambda]$. Then $[\lambda]$ is supported by L . Furthermore L does not depend on the choice of $\{c_n\}$, as long as $\{c_n^*\}$ exits every compact subset of E .

Combining the end invariants of the geometrically finite ends and the ending laminations of the geometrically infinite ends, we get the end invariants of ρ . In section 2.8, we describe more precisely the kind of objects thus obtained.

2.6 Thurston's compactification

The Teichmüller space $\mathcal{T}(S)$ of a hyperbolic surface S can be compactified by actions on $\mathbb{R}$ -trees, up to homothety. The actions thus obtained are small and minimal. By Skora's Theorem, see [Sk], such an action is dual to a measured geodesic lamination. Since $\mathcal{T}(S)$ is compactified by actions on $\mathbb{R}$ -trees up to homothety, we actually get a projective measured geodesic lamination. In order to prove some results in the present paper, we need a more precise description of the behavior of a sequence $\{m_n\} \subset \mathcal{T}(S)$ tending to a projective measured geodesic lamination $[\lambda]$. In this section we recall how Thurston compactifies $\mathcal{T}(S)$ directly by projective measured geodesic laminations (that is, without using $\mathbb{R}$ -trees), following the exposition given in [FLP, Exposé 8].

Let m be a complete hyperbolic metric on S . Let $P \subset S$ be a multi-curve such that the components of $S - P$ are 3-holed spheres. We refer to a 3-holed sphere as a *pair of pants*, and such a multi-curve P is called a *pants decomposition* of S . Following [FLP, §I.2. Exposé 8] and using P , we associate a partial *measured foliation* $\mathcal{F}_P(m)$ of S to m . (Here, a *measured foliation* of a surface S is a singular foliation of S with a transverse measure, where the measure of a curve transverse to the foliation is preserved by homotopies preserving the foliation. A *partial measured foliation* is a measured foliation of a subset of S .) Let R be a pair of pants and let l_1, l_2, l_3 be the leaves of P bounding R .

Assume first that $\ell_m(l_i) \leq \ell_m(l_j) + \ell_m(l_k)$ for any $i \neq j \neq k$. For $i \neq j$, let $k_{i,j}$ be the geodesic segment orthogonal to l_i and l_j . Let $T_{i,j}$ be the set of points at a distance at most $\frac{1}{4}(\ell_m(l_i) + \ell_m(l_j) - \ell_m(l_k))$ from $k_{i,j}$. This set is foliated by the curves $\{z \mid d(k_{i,j}, z) \text{ is constant}\}$ and the transverse measure is given by the distance between 2 leaves. Thus in R we have three foliated sets. Notice that by the choice of their widths they don't intersect in $\text{int}(R)$. Thus we have constructed a partial measured foliation $\mathcal{F}_P(m)$ of R (see figure 1).




Figure 1: The partial foliation of R

If on the other hand $\ell_m(l_1) > \ell_m(l_2) + \ell_m(l_3)$ (up to relabeling, this is the only case left to consider) we take $T_{1,2}$, respectively $T_{1,3}$, to be the set of points at a distance at most $\frac{\ell_m(l_2)}{2}$ from $k_{1,2}$, foliated as before, respectively the set of points at a distance at most $\frac{\ell_m(l_3)}{2}$ from $k_{1,3}$. Let $k_{1,1}$ be the (simple) geodesic segment orthogonal at both ends to l_1 . The set of points of l_1 not lying in the interior of $T_{1,2} \cap l_1$ nor of $T_{1,3} \cap l_1$ form 2 arcs A and A' . We take $T_{1,1}$ to be the union of the curves $\{z \mid d(k_{1,1}, z) \text{ is constant}\}$ starting at A (notice that $k_{1,1}$ may not lie in $T_{1,1}$). Thus we have constructed a partial measured foliation $\mathcal{F}_P(m)$ of R (see figure 1).

It is proved in [FLP, Exposé 8] that the projection mapping m to $\mathcal{F}_P(m)$ is a homeomorphism from $\mathcal{T}(S)$ into the set of partial measured foliations giving a non-zero measure to any leaf of P . Notice that $\mathcal{F}_P(m)$ depends on the choice of P .

One can extend this measured partial foliation into a singular measured foliation of the whole surface. This singular foliation is well defined up to isotopy. It has been noticed by Thurston that there is a natural bijection between equivalence classes (up to isotopy and Whitehead moves, see [FLP]) of singular measured foliations of a surface and measured geodesic laminations, see [Lev]. To a measured foliation $\mathcal{F}_P(m)$, there corresponds the measured geodesic lamination λ satisfying $i(\mathcal{F}_P(m), c) = i(\lambda, c)$ for any simple closed curve c . From now on we give the same name to a foliation and to the corresponding lamination.

The measured geodesic lamination $\mathcal{F}_P(m)$ roughly describes the length spectrum of m , as can be seen in the following result, see [Th3, Theorem 2.2]:

Theorem 2.2. *Let m_0 be a complete hyperbolic metric on S , and let P be a pants decomposition of S . Let $\varepsilon > 0$ be such that $\ell_{m_0}(c) \geq \varepsilon$ for any leaf c of P . Then there is $Q = Q(\varepsilon)$ such that any measured geodesic lamination λ on S satisfies $i(\lambda, \mathcal{F}_P(m_0)) \leq \ell_{m_0}(\lambda) \leq i(\lambda, \mathcal{F}_P(m_0)) + Q\ell_{m_0}(\lambda)$. $\square$*

Now we explain how a projective measured geodesic lamination is associated to a diverging sequence of points of $\mathcal{T}(S)$. First, note that there exists $\varepsilon > 0$ so that if a simple closed geodesic $c \subset S$ has length less than ε then any simple closed geodesic crossing c has length at least ε . The existence of such an ε can be easily deduced from the Collar Lemma and ε does not depend on the hyperbolic metric on S . Take any pants decomposition Q of S . Given a sequence $\{m_n\} \subset \mathcal{T}(S)$, we extract a subsequence such that there exists N such that for any leaf c of Q either $\ell_{m_n}(c) \leq \varepsilon$ for any $n \geq N$ or $\ell_{m_n}(c) \geq \varepsilon$ for any $n \geq N$. If a leaf c of Q satisfies $\ell_{m_n}(c) \leq \varepsilon$ for any $n \geq N$ then we replace it by a simple closed curve d that crosses c and is disjoint from $Q - c$. Thus we get a new pants decomposition P such that for any leaf c of P , $\ell_{m_n}(c) \geq \varepsilon$ for any $n \geq N$. Now we consider $\mathcal{F}_P(m_n)$. Assume that $\{m_n\}$ is a diverging sequence. By Theorem 2.2, $\mathcal{F}_P(m_n)$ diverges as well. We extract a subsequence so that $\{[\mathcal{F}_P(m_n)]\}$ converges in the space of projective measured geodesic laminations.

Thus we get Thurston's compactification of Teichmüller space by projective measured geodesic laminations. Using Theorem 2.2 it is easy to see that this compactification is the same as the one obtained by combining the Morgan-Shalen compactification by actions on $\mathbb{R}$ -trees with Skora's Theorem.

Forgetting the transverse measure, we say that a sequence $\{m_n\}$ of complete hyperbolic metrics on S tends to a lamination L on S if any subsequence contains a further subsequence

converging to a projective measured geodesic lamination supported by L .

2.7 Compactification of Teichmüller space for surfaces with boundary

Given a sequence of complete hyperbolic metrics $\{m_n\}$ on a closed surface, we will need to describe the behavior of the sequence on some specified subsurfaces. Namely we need to define a compactification of Teichmüller space for surfaces with boundary.

The fastest way to do so is to use the compactification by actions on $\mathbb{R}$ -trees and Skora's Theorem. Consider a compact surface F with boundary and let $\{m_n\}$ be a sequence of hyperbolic metrics on F such that ∂F is an union of m_n -geodesics for any n . Assume that $\{m_n\}$ does not contain a convergent subsequence and that there is a non-peripheral simple closed curve $c \subset F$ such that $\frac{\ell_{m_n}(\partial F)}{\ell_{m_n}(c)} \rightarrow 0$. We use c to make sure that there is a non peripheral closed curve whose length grows much faster than the length of ∂F . This ensures that each element of ∂F , viewed as a conjugacy class in $\pi_1(S)$, has a fixed point when acting on the $\mathbb{R}$ -tree to which $\{m_n\}$ tends. Such a fixed point is necessary to use Culler-Morgan-Shalen's Theory and Skora's Theorem. Then, by [MoS1] and [Sk], a subsequence of $\{m_n\}$ tends to an action of $\pi_1(F)$ on an $\mathbb{R}$ -tree dual to a compact geodesic measured geodesic lamination λ .

For technical reasons, we will need to associate to a metric m on F a measured geodesic lamination, as was done for closed surfaces. Let m be a hyperbolic metric on F such that ∂F is an union of closed geodesics. Let P_F be a pants decomposition of F , so that the connected components of $F - (P_F \cup \partial F)$ are three-holed spheres. (Hence, we do not consider ∂F to be contained in P_F .) Fix $\varepsilon > 0$ so that $\ell_m(c) \geq \varepsilon$ for any leaf c of P_F . We are especially interested in sequences of metrics on F with arbitrarily short boundary curves. In particular, we cannot simply repeat the construction of $\mathcal{F}_P(m)$ in each component of $F - P_F$ as we did in the case of a closed surface, as we won't then satisfy the hypothesis of Theorem 2.2.

Consider the surface DF obtained by doubling F along its boundary, i.e. DF is obtained by taking F and its mirror image and by identifying the corresponding boundaries of these two surfaces. Endowing $F \subset DF$ with a hyperbolic metric m with geodesic boundary and its mirror image with the mirror image of m , we get a complete hyperbolic metric Dm on DF . We denote by $\partial F \subset DF$ the multi-curve corresponding to the identified boundaries of F and its mirror image. We consider the pants decomposition P_F and its mirror image; the union yields a multi-curve $DP_F \subset DF$. There is a natural involution $\tau : DF \rightarrow DF$ that exchanges F with its mirror image. By construction we have $\tau(DP_F) = DP_F$. We complete DP_F into a pants decomposition P_{DF} so that we have $\tau(P_{DF}) = P_{DF}$ (see figure 2). There are 2 possibilities for a component of $DF - DP_F$ that intersects ∂F , and figure 2 shows how to extend DP_F in both cases.

Now we consider the map $\mathcal{F}_{P_{DF}} : \mathcal{T}(DF) \rightarrow \mathcal{ML}(DF)$ associated to P_{DF} as defined in section 2.6. When we have $i(\mathcal{F}_{P_{DF}}(Dm), \partial F) = 0$, then the restriction of $\mathcal{F}_{P_{DF}}(m)$ to the copy of F used to construct DF is a measured geodesic lamination $\mathcal{F}_{P_{DF}}(m)$. The metric m is uniquely defined by $\mathcal{F}_{P_{DF}}(m)$ and the lengths of the components of ∂F . Thus we have a well-defined injective map from the set of metrics on F satisfying $i(\mathcal{F}_{P_{DF}}(Dm), \partial F) = 0$ to $\mathcal{ML}(\text{int}(F)) \times \mathbb{R}_+^k$ (where k is the number of components of ∂F).




Figure 2: Extending DP_F to a pants decomposition P_{DF}

Let us show that when $\ell_m(\partial F)$ is small enough, we have $i(\mathcal{F}_{P_{DF}}(Dm), \partial F) = 0$.

Claim 2.3. *Given $\varepsilon > 0$, there exists $\eta > 0$ depending only on ε such that if $\ell_m(\partial F) \leq \eta$ then $i(\mathcal{F}_{P_{DF}}(Dm), \partial F) = 0$.*

Proof. Consider the partial measured foliation $\mathcal{F}_{P_{DF}}(Dm)$ as originally defined and let $|\mathcal{F}_{P_{DF}}(Dm)|$ be its support. By [FLP, Exposé 8], there is a uniform bound K on the length of the closure of any component of $|\mathcal{F}_{P_{DF}}(Dm)| \cap (DF - P_{DF})$, depending only on ε . By the Margulis Lemma, we can find η so that any component of DP_F is at distance at least $K + 1$ from ∂F . With this choice of η , the closure of a component of $|\mathcal{F}_{P_{DF}}(Dm)| \cap (DF - P_{DF})$ with one endpoint on DP_F cannot intersect ∂F .

Since P_{DF} is invariant under the action of τ , we have $\tau(\mathcal{F}_{P_{DF}}(Dm)) = \mathcal{F}_{P_{DF}}(Dm)$. Furthermore ∂F is fixed pointwise by the action of τ . It follows that the closure of a component of $|\mathcal{F}_{P_{DF}}(Dm)| \cap (DF - P_{DF})$ intersects ∂F only if it has both endpoints in DP_F .

Combining these two paragraphs, we conclude that for η small enough no leaf of $\mathcal{F}_{P_{DF}}(Dm)$ crosses ∂F . In particular, we have $i(\mathcal{F}_{P_{DF}}(Dm), \partial F) = 0$. $\square$

Given a sequence of metrics $\{m_n\}$ on F such that $\ell_{m_n}(\partial F) \rightarrow 0$, we choose a pants decomposition P_F (not containing ∂F) so that we have $\ell_{m_n}(c) \geq \varepsilon$ for any leaf c of P_F and any n . We construct a map $\mathcal{F}_{P_{DF}} : \{m_n\} \rightarrow \mathcal{ML}(F)$ as described previously. Taking a subsequence $\{m_n\}$ such that $\mathcal{F}_{P_{DF}}(m_n)$ converges projectively, we get a projective measured geodesic lamination $[\lambda]$. Since Theorem 2.2 holds for Dm_n and $\mathcal{F}_{P_{DF}}(Dm_n)$, it is still true when applied to m_n and $\mathcal{F}_{P_{DF}}(m_n)$. It follows that $\{m_n\}$ tends to an action on an $\mathbb{R}$ -tree which is dual to $[\lambda]$.

A complete metric m on an open surface F can be approximated by a sequence of metrics m_n on the surface $\overline{F}$ obtained by adding a simple closed curve along each cusp of F . The metrics m_n have the property that $\partial \overline{F}$ is a union of m_n -geodesics whose lengths tend to 0. Choosing an appropriate pants decomposition P of F (again not containing ∂F), one can define measured geodesic laminations $\mathcal{F}_{P_{DF}}(m_n)$ associated to $\{m_n\}$ as above. It is easy to see that $\{\mathcal{F}_{P_{DF}}(m_n)\}$ converges to a measured geodesic lamination $\mathcal{F}_{P_{DF}}(m)$. Notice that $\mathcal{F}_{P_{DF}}(m)$ depends on P but not on the choice of the sequence $\{m_n\}$. We have thus defined a map $\mathcal{F}_{P_{DF}} : \mathcal{T}(F) \rightarrow \mathcal{ML}(F)$ for which Theorem 2.2 holds.

2.8 Gallimaufries

Let S be a compact orientable surface of genus at least 2, not necessarily connected. A *gallimaufry* $\Gamma = (F, m, L)$ on S is made as follows: F is an open incompressible subsurface of S , namely its closure is an incompressible subsurface, and in particular a connected component of F is either a surface with cusps or a connected component of S ; m is a complete hyperbolic metric (up to isotopy) on F with finite area; and L is a recurrent geodesic lamination on the closed surface $S - F$. We furthermore require that L does not have any closed leaf and that the connected components of $S - (F \cup L)$ are discs, annuli or Möbius bands. The surface F is the *moderate surface* of Γ and the lamination L forms its *immoderate lamination*. We denote by $\mathcal{GA}(S)$ the set of all gallimaufries on S . As an example, the end invariants of a hyperbolic 3-manifold form a gallimaufry: the union of the open surfaces facing the geometrically finite ends forms its moderate surface, equipped with the metric induced by the corresponding conformal structure at infinity; the union of the ending laminations of its geometrically infinite ends is its immoderate lamination. We now define the topology on $\mathcal{GA}(S)$.

Let $\{\Gamma_n = (F_n, m_n, L_n)\}$ be a sequence of gallimaufries on S . For each component of $S - F_n$ which is an annulus or Möbius band A , we add to L_n an essential simple closed curve that can be homotoped into A . Thus we get a new geodesic lamination L'_n . We say that $\{\Gamma_n\}$ converges to $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$ in $\mathcal{GA}(S)$ if the following hold:

- i) for any n we have $F_\infty \subset F_n$ and the restrictions of the m_n to F_∞ converge to m_∞ , namely we have $l_{m_n}(\partial \overline{F}_\infty) \rightarrow 0$ and for any non-peripheral closed curve $c \subset F_\infty$, we have $\ell_{m_n}(c) \rightarrow \ell_{m_\infty}(c)$,
- ii) the recurrent part of the Hausdorff limit of any convergent subsequence of $\{L'_n\}$ lies in L'_∞ ,
- iii) if a component L of L_∞ lies in infinitely many F_n then the restrictions of the m_n to $S(L)$ tend to L .

A gallimaufry with empty immoderate lamination is simply a point in the Teichmüller space of S . Thus by defining a gallimaufry we have constructed a bordification of Teichmüller space. Notice that $\mathcal{GA}(S)$ is not compact and not even locally compact.

Lemma 2.4. *The space $\mathcal{GA}(S)$ is not locally compact.*

Proof. Consider a simple closed curve $c \subset S$ and an embedded annulus $A \subset S$ around c . Let m be a complete hyperbolic metric on the open surface $S - A$ and let $\{m_n\}$ be a sequence of complete hyperbolic metrics on S converging to m on $S - A$ in the sense of i) above. In particular since c corresponds to a cusp of $S - A$, we have $l_{m_n}(c) \rightarrow 0$.

Let $\phi : S \rightarrow S$ be a (right) Dehn twist along c . By construction the sequence of gallimaufries $\{(S, m_n, \emptyset)\}$ converges to $(S - A, m, \emptyset)$ and since we have $\ell_{m_n}(c) \rightarrow 0$, for any fixed k , the sequence $\{(S, \phi_*^k m_n, \emptyset)\}$ also converges to $(S - A, m, \emptyset)$. On the other hand if we fix n and let $k \rightarrow \infty$, no subsequence of $\{(S, \phi_*^k m_n, \emptyset)\}$ converges in $\mathcal{GA}(S)$ for the following reason. First notice that if a sequence of metrics $\{m_n\}$ converges to a gallimaufry

$\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$, it follows from the definitions that we have $l_{m_n}(\partial F_\infty) \rightarrow 0$. On the other hand, the $\phi_*^k m_n$ -length of any given simple closed curve is bounded away from 0, when n is fixed and k varies. It follows that if $\{(S, \phi_*^k m_n, \emptyset)\}$ converges to a gallimaufry $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$ then we would have $F_\infty = S$ and $L_\infty = \emptyset$. This would mean that the metrics $\{\phi_*^k m_n\}$ converge, when n is fixed and k varies, which is clearly not the case.

Thus for any given compact set K , we can find a sequence $\{k_n\}$ so that the sequence $\{(S, \phi_*^{k_n} m_n, \emptyset)\}$ eventually exits K . On the other hand, the sequence $\{(S, \phi_*^{k_n} m_n, \emptyset)\}$ converges to $(S - A, m, \emptyset)$. Hence we have shown that $\mathcal{GA}(S)$ is not locally compact. $\square$

2.9 Double incompressibility

Let M be a compact 3 manifold and let $F \subset \partial M$ be an incompressible surface. We say that a geodesic lamination $L \subset F$ is *doubly incompressible in F* if it contains the support of a measured geodesic lamination λ satisfying the following condition: there exists $\eta > 0$ such that for any essential disc, annulus or Möbius band $(E, \partial E)$ properly embedded in (M, F) , we have $i(\lambda, \partial E) \leq \eta$.

Notice that if a measured geodesic lamination L is doubly incompressible, then any measured geodesic lamination with support $|L|$ is also doubly incompressible. Thus it only depends on the topology of L .

Say that a gallimaufry $\Gamma = (F, m, L)$ is *doubly incompressible* if L is doubly incompressible in $\partial M - F$. The end invariants of a compact 3 manifold form a doubly incompressible gallimaufry, see [Bon] and [Ca].

3 Algebraic convergence

In this section we prove that the convergence of the end invariants to a doubly incompressible gallimaufry implies the algebraic convergence of the corresponding representations (up to taking a subsequence).

Proposition 3.1. *Let $\{\rho_n\} \subset \text{AH}(M)$ be a sequence of representations uniformizing M and let $\Gamma_n = (F_n, m_n, L_n)$ be the end invariants of $\rho_n(\pi_1(M))$. Assume that $\{\Gamma_n\}$ converges in $\mathcal{GA}(\partial M)$ to a doubly incompressible gallimaufry $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$. Then a subsequence of $\{\rho_n\}$ converges algebraically.*

We prove this proposition by contradiction, roughly following the plan of Otal's proof of the Double Limit Theorem, see [Ot2]. Assume that no subsequence of $\{\rho_n\}$ converges algebraically. By work of Morgan and Shalen, see [MoS1], a subsequence of $\{\rho_n\}$ tends to a small minimal action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$. Consider a component L of the immoderate lamination L_∞ of Γ_∞ . Using the assumption that $\{\Gamma_n\}$ converges to Γ_∞ we construct a sequence of laminations sufficiently close to L whose lengths are sufficiently well controlled. We then deduce from the work of Otal, see [Ot1], that L cannot be realized in $\mathcal{T}$. On the other hand we see that since Γ_∞ is doubly incompressible, at least one component of its immoderate lamination is realized in $\mathcal{T}$. This yields the expected contradiction.

3.1 Cut and paste

Before starting the proof of Proposition 3.1, we will describe a relatively straightforward cut and paste operation that will be used many times throughout this paper.

Consider a closed orientable surface S and an essential open subsurface $F \subset S$ which is not a pair of pants. Let $c \subset S$ be a simple closed curve that intersects F . We want to construct a simple closed curve $e \subset F$ that behaves somewhat like c . If c lies in F then we take $e = c$. Otherwise, let k be a component of $c \cap \overline{F}$; it is an arc joining two boundary components of $\overline{F}$ (which may not be distinct). Let $\mathcal{V}$ be a small neighborhood of the union of k and of the components of $\partial\overline{F}$ containing the endpoints of k . The boundary of $\mathcal{V} \cap F$ contains 1 or 2 simple closed curves depending on whether the endpoints of k lie in different components of $\partial\overline{F}$. Since F is not a pair of pants, at least one of these curves is not peripheral. Let e be such a simple closed curve, namely e is freely homotopic to a component of $\partial\mathcal{V}$ and is not peripheral in $\overline{F}$.

By construction we have:

Claim 3.2. *The simple closed curve e satisfies the inequalities $\ell_s(e) \leq 2\ell_s(c) + \ell_s(\partial\overline{F})$ and $i(e, c) \leq i(c, \partial\overline{F})$.* $\square$

Consider now a sequence of simple closed curves $\{c_n\} \subset S$ and the sequence of simple closed curves $\{e_n\} \subset F$ produced by the operation above. Extract subsequences so that $\{c_n\}$ and $\{e_n\}$ converge in the Hausdorff topology to geodesic laminations C and E respectively. Assuming that F contains a minimal sublamination L of C , we have:

Claim 3.3. *The lamination L is a sublamination of E .*

Proof. Put a weight u_n on c_n so that $\{u_n c_n \cap F\}$ converges to a measured geodesic lamination λ supported by L . With such weights, we have $i(u_n c_n, \partial\overline{F}) \rightarrow 0$. Since we have $i(e_n, c_n) \leq i(c_n, \partial\overline{F})$ we get $i(u_n c_n, e_n) \rightarrow 0$. Put a weight v_n on e_n so that $\{v_n e_n\}$ converges to a measured geodesic lamination μ . The support of μ is a sublamination of E . Since $\{v_n\}$ is bounded, we have $i(\lambda, \mu) = \lim i(u_n c_n, v_n e_n) \rightarrow 0$. Hence λ and μ have the same support and L is a sublamination of E . $\square$

Using this Claim, we will describe the behavior of a sequence of simple closed curves with bounded length in a sequence of metrics that degenerates.

Lemma 3.4. *Let $\{m_n\}$ be a sequence of metrics on a closed surface S . Let $F \subset S$ be an essential subsurface such that the restrictions of the $\{m_n\}$ to F converge to a geodesic lamination L (in the sense of Section 2.7). Let $\{c_n\}$ be a sequence of simple closed curves on S such that $\{\ell_{m_n}(c_n)\}$ is a bounded sequence and that $\{c_n\}$ intersects $S(L)$ for n large enough. Extract a subsequence such that $\{c_n\}$ converges in the Hausdorff topology to a geodesic lamination C . Then L is a sublamination of C .*

Proof. Using the cut and paste construction described above, we get a sequence of simple closed curves $\{e_n \subset S(L)\}$ with $\ell_{m_n}(e) \leq 2\ell_{m_n}(c) + \ell_{m_n}(\partial S(L))$ and $i(e, c) \leq i(c, \partial S(L))$. Consider a sequence $\{u_n\}$ converging to 0 such that the sequence $\{u_n e_n\} \subset \mathcal{ML}(S(L))$ converges to a measured geodesic lamination λ . By Claim 3.3, the support of λ is a sublamination

of C .

Consider the double $\overline{DS(L)}$ of $\overline{S(L)}$ as defined in Section 2.7 and the metric Dm_n on $\overline{DS(L)}$ induced by the restriction of m_n to $\overline{S(L)}$. Choose a pants decomposition P of $\overline{DS(L)}$ as in Section 2.7 such that there is $\varepsilon > 0$ for which we have that $\ell_{Dm_n}(d) \geq \varepsilon$ for any n and any leaf d of P and that P is invariant under the natural involution τ of $\overline{DS(L)}$. We have a measured geodesic lamination $\mathcal{F}_p(Dm_n)$ defined as in Section 2.6.

Extract a subsequence such that $\{|\mathcal{F}_p(Dm_n)|\}$ converges in the Hausdorff topology to a geodesic lamination L' . Then we have $\tau(L') = L'$. In particular if a component d of $\partial\overline{S(L)}$ is a leaf of L' then it is an isolated leaf. Let us remove from $\mathcal{F}_p(Dm_n)$ any leaf that is a component of $\partial\overline{S(L)}$. Since the restrictions of the m_n to $S(L)$ tends to L , up to extracting a subsequence there are $v_n \rightarrow 0$ such that $\{v_n \mathcal{F}_p(Dm_n)\}$ converges to a measured geodesic lamination $D\mu$ whose support is the “double” of L . By definition, we have $v_n l_{Dm_n}(\partial\overline{S(L)}) \rightarrow i(D\mu, \partial\overline{S(L)}) = 0$. We denote by e_n the curve defined by e_n on the copy of $\overline{S(L)}$ comprising $\overline{DS(L)}$. From Theorem 2.2, we get $i(e_n, v_n \mathcal{F}_p(Dm_n)) \leq v_n l_{m_n}(c_n) + 2v_n l_{m_n}(\partial\overline{S(L)}) \rightarrow 0$. It follows that $i(\lambda, D\mu) = \lim u_n v_n i(e_n, \mathcal{F}_p(Dm_n)) = 0$. Hence the support of λ is L . Now from the first paragraph, we can conclude that L is a sublamination of C . $\square$

3.2 Length and realization

Consider a component L of the immoderate lamination L_∞ of Γ_∞ . In this section, we show that L cannot be realized in $\mathcal{T}$. To do that, we construct a sequence of geodesic laminations with controlled length which are close enough (in a sense to be made precise) to L . We start by roughly approximating L by simple closed curves of bounded length, using Lemma 3.4.

Claim 3.5. *There is a sequence $\{c_n\}$ of simple closed curves on S such that $\{\ell_{\rho_n}(c_n^*)\}$ is bounded and so that, up to extracting a subsequence, $\{c_n\}$ converges in the Hausdorff topology to a geodesic lamination containing L .*

Proof. We first assume that $L \subset F_n$ for n sufficiently large, where F_n is the moderate surface of Γ_n . It is a classical result of Bers [Be] that there are pants decompositions P_n of the F_n so that $\{\ell_{m_n}(P_n)\}$ is a bounded sequence. Extract a subsequence such that $\{P_n\}$ converges in the Hausdorff topology to a geodesic lamination P_∞ . Since P_n is a pants decomposition of F_n and since F is an essential subsurface of F_n , for any n there is a leaf c_n of P_n that intersects F . By assumption, $\ell_{m_n}(c_n)$ is bounded. Furthermore $\{c_n\}$ converges in the Hausdorff topology to a sublamination C of P_∞ . By Lemma 3.4, L is a sublamination of C . By [Ah], we have $\ell_{\rho_n}(c_n^*) \leq 2\ell_{m_n}(c_n)$. In particular $\{\ell_{\rho_n}(c_n^*)\}$ is bounded.

If we have $L \not\subset F_n$ then L lies in the Hausdorff limit of any convergent subsequence of $\{L'_n\}$ where L'_n is obtained by adding to L_n a curve in each component of $\partial_{X<0}M - F_n$ which is an annulus or a Möbius band. By the definition of L'_n , if a component B_n of L'_n intersecting $S(L)$ is not a closed leaf, it is an ending lamination of ρ_n . In this case, there are curves c_k such that $\ell_{\rho_n}(c_k^*) \leq Q$ for some Q depending only on ∂M , such that $\{c_k\}$ converges in the Hausdorff topology to a geodesic lamination containing B_n . Taking a diagonal sequence we get a sequence $\{c_n\}$ of simple closed curves so that the sequence $\{\ell_{\rho_n}(c_n^*)\}$ is bounded and so that $\{c_n\}$ converges in the Hausdorff topology to a geodesic lamination containing L . If

B_n is a closed curve, then $\ell_{\rho_n}(B_n^*) = 0$ and we are done by taking $c_n = B_n$. This concludes the proof of Claim 3.5. $\square$

Now that we have this sequence $\{c_n\}$, we use the following proposition to deduce that no component of L is realized in $\mathcal{T}$.

Lemma 3.6. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\{\rho_n\} \subset \text{AH}(M)$ be a sequence tending to a small minimal action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$. Let $\{c_n\} \subset \partial M$ be a sequence of simple closed curves such that $\{\ell_{\rho_n}(c_n^*)\}$ is bounded. Assume that $\{c_n\}$ converges in the Hausdorff topology to a geodesic lamination C_∞ and let C be an irrational minimal sublamination of C_∞ . Then C is not realized in $\mathcal{T}$.*

Proof. The study of the behavior of the lengths of geodesic laminations that are realized in $\mathcal{T}$ has already been initiated by J.-P. Otal, see [Ota1]. His results are stated under the assumption that M is a handlebody, and he considers a sequence of geodesic laminations that converges in the Hausdorff topology. But a careful look at the proof yields the following statement.

Theorem 3.7 (Continuity Theorem [Ota1]). *Let M be a compact, orientable, hyperbolizable 3-manifold. Let $\{\rho_n\}$ be a sequence of geometrically finite representations of $\pi_1(M)$ tending to a small minimal action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$. Let $\varepsilon_n \rightarrow 0$ be such that for all $g \in \pi_1(M)$, we have $\varepsilon_n \ell_{\rho_n}(g) \rightarrow \ell_{\mathcal{T}}(g)$ and let $L \subset \partial_{\chi < 0} M$ be a minimal geodesic lamination which is realized in $\mathcal{T}$. Consider a geodesic lamination $E \subset S(L)$ containing L . Then there exists a neighborhood $\mathcal{V}(E)$ of E and constants Q, n_0 such that for any simple closed curve $c \subset \mathcal{V}(L)$ and for any $n \geq n_0$,*

$$\varepsilon_n \ell_{\rho_n}(c^*) \geq Q \ell_{s_0}(c).$$

In this statement, s_0 is a reference metric which is used to measure the "complexity" of the curve c . Any complete hyperbolic metric on ∂M can be chosen and Q will depend on this choice.

Theorem 3.7 is enough to conclude the proof of Lemma 3.6 when $\{c_n\}$ converges in the Hausdorff topology to L . In order to deal with the more general case, we use the cut and paste operation described in section 3.1 in $\overline{S(L)}$ on the c_n . This provides us with a sequence of simple closed curves $\{e_n\} \subset S(L)$ satisfying: $\ell_s(e_n) \leq 2\ell_s(c) + \ell_s(\partial \overline{F})$ and $i(e_n, c) \leq i(c, \partial \overline{F})$. Furthermore, by Lemma 3.3, up to extracting a subsequence, $\{e_n\}$ converges in the Hausdorff topology to a geodesic lamination E containing L .

Now it remains to control the length in $M_n = \mathbb{H}^3 / \rho_n(\pi_1(M))$ of the sequence $\{e_n\}$ thus constructed. Let S be the connected component of ∂M containing L . Denote by $\phi : S \rightarrow S$ the mapping class that performs one left Dehn twist in each component of $W(L) - S(L)$. For a fixed n , the sequence $\{\phi^k(c_n)\}$ converges as $k \rightarrow \infty$ to a finite geodesic lamination whose non-compact leaves spiral in $W(L) - S(L)$. Extend this lamination to a finite lamination T_n whose non-compact leaves spiral in $W(L) - S(L)$ and whose complementary regions are ideal triangles. Let $f_n : S \rightarrow M_n$ be a pleated surface (see definition in [CEG, §5.1]) homotopic to the inclusion map such that f_n maps any leaf of T_n to a geodesic of M_n . The existence of such a pleated surface follows from [CEG, §5.3]. We denote by $\ell_{f_n}(d)$ the length of a closed geodesic d of S endowed with the metric induced by f_n . If a component c of $\partial S(L)$

corresponds to a parabolic isometry in $\rho_n(\pi_1(M))$, we consider a map $f_n : S - c \rightarrow M_n$ such that the cusps of $S - c$ are mapped to the corresponding cusps of M_n . Such a map f_n is called a *noded pleated surface*, see [Mi1], and we set $\ell_{f_n}(c) = 0$ in this case.

Given $\varepsilon > 0$, it follows from the "efficiency of pleated surfaces", see [Th3, Theorem 3.3], that there is a constant $Q = Q(\varepsilon)$ such that $\ell_{f_n}(c_n \cap R_D) \leq \ell_{\rho_n}(c_n^*) + Q i(c_n, E)$, where R_D is the complement of the ε -Margulis tubes around the components of $\partial S(L)$ with short length (with respect to the metric induced by f_n). (Compare with [Mi1, p. 138].) It follows that there is a component k_n of $c_n \cap S(L) \cap R_D$ such that $\{\ell_{f_n}(k_n)\}$ is bounded by $K > 0$. Using these arcs k_n in the construction of Lemma 3.1, we get a lamination $E \subset S(L)$ with $L \subset E$ and a sequence of simple closed curves $\{e_n\} \subset \mathcal{ML}(S(L))$ such that $\{e_n\}$ converges to E in the Hausdorff topology and such that $\ell_{f_n}(e_n) \leq 2\ell_{f_n}(k_n) + \ell_{f_n}(\partial \overline{S(L)})$.

By the choice of k_n , the sequence $\{\ell_{f_n}(k_n)\}$ is bounded by K , and so $\ell_{f_n}(e_n) \leq 2K + \ell_{f_n}(\partial \overline{S(L)})$. Notice that since f_n realizes $\partial \overline{S(L)}$, we have $\ell_{f_n}(\partial \overline{S(L)}) = \ell_{\rho_n}(\partial \overline{S(L)}^*)$. Thus we get $\ell_{\rho_n}(e_n^*) \leq 2K + \ell_{\rho_n}(\partial \overline{S(L)}^*)$. Since the action of $\rho_n(\pi_1(M))$ tends to the action of $\pi_1(M)$ on the $\mathbb{R}$ -tree $\mathcal{T}$, there is a sequence $\varepsilon_n \rightarrow 0$ such that we have $\varepsilon_n \ell_{\rho_n}(g) \rightarrow \ell_{\mathcal{T}}(g)$ for any $g \in \pi_1(M)$. In particular, we have $\varepsilon_n \ell_{\rho_n}(\partial \overline{S(L)}^*) \rightarrow \ell_{\mathcal{T}}(\partial \overline{S(L)})$. Thus $\varepsilon_n \ell_{\rho_n}(e_n^*) \leq 2\varepsilon_n K + \varepsilon_n \ell_{\rho_n}(\partial \overline{S(L)}^*)$ is bounded.

On the other hand, since $\{e_n\}$ converges to $E \supset L$, we have $\ell_{s_0}(e_n) \rightarrow \infty$ for any complete hyperbolic metric s_0 on S . It follows then from Theorem 3.7 that $C \subset E$ is not realized in $\mathcal{T}$. This concludes the proof of Lemma 3.6. $\square$

Combining Claim 3.5 and Lemma 3.6, we conclude that no component of L_∞ is realized in $\mathcal{T}$.

3.3 Double incompressibility and realization

Now we show that when Γ_∞ is doubly incompressible, at least one minimal sublamination of L_∞ is realized. Thus we get a contradiction with Claim 3.5 and Lemma 3.6.

Lemma 3.8. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Consider a sequence of representations $\{\rho_n\} \subset \text{AH}(M)$ uniformizing M that tends to a small minimal action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$. Let $\Gamma = (F, m, L)$ be a doubly incompressible gallimaufry. Assume that for any simple closed curve $c \subset F$, the sequence $\{\ell_{\rho_n}(c^*)\}$ is bounded. Then at least one component of L is realized in $\mathcal{T}$.*

Proof. Consider a component S of $\partial_{\chi < 0} M$. Since ∂M is incompressible, we can view $\pi_1(S)$ as a subgroup of $\pi_1(M)$. Thus we have a small action of $\pi_1(S)$ on $\mathcal{T}$. Let $\mathcal{T}_S$ be the minimal sub-tree of $\mathcal{T}$ for this action. By Skora's Theorem, see [Sk], this action is dual to a measured geodesic lamination β . Doing the same for each component of ∂M we get a measured geodesic lamination $\beta \in \mathcal{ML}(\partial M)$. By [MoS2], β is not trivial. By [Ot1], if a minimal geodesic lamination crosses $|\beta|$, then it is realized in $\mathcal{T}$. So we have to show that one component of L crosses $|\beta|$.

For a simple closed curve $c \subset F$, we have assumed that $\{\ell_{\rho_n}(c^*)\}$ is bounded. Hence we have $i(c, \beta) = 0$ for any simple closed curve $c \subset F$. This is possible only if $|\beta|$ lies in $\partial M - F$. By [Le1, Lemme 4.7], see also [BoO], there is a sequence $\{E_n\}$ of essential annuli or Möbius

bands such that $\{\partial E_n\}$ converges in the Hausdorff topology to a sublamination B of $|\beta|$. Since L is doubly incompressible in $(M, \partial M - F)$, L crosses B . Thus we have proved that $|\beta|$ crosses L . By [Ot1, Theorem 3.1.4] this concludes the proof of Lemma 3.8 $\square$

We can now conclude the proof of Proposition 3.1.

Proof of Proposition 3.1. Assume that the conclusion is not satisfied, namely that a subsequence of $\{\rho_n\}$ tends to an action of $\pi_1(M)$ on an $\mathbb{R}$ -tree $\mathcal{T}$. By Claim 3.5 and Lemma 3.6 no component of L_∞ is realized in $\mathcal{T}$. By the definition of the topology on $\mathcal{GA}(\partial M)$, for any simple closed curve $c \subset F_\infty$, the sequence $\{\ell_{m_n}(c)\}$ is bounded. By [Ah], the sequence $\{\ell_{\rho_n}(c^*)\}$ is also bounded. Thus the hypotheses of Lemma 3.8 are fulfilled. It follows that at least one component of L_∞ is realized in $\mathcal{T}$. We conclude from this contradiction that a subsequence of $\{\rho_n\}$ converges algebraically (up to conjugacy). $\square$

4 Strong convergence

Let M be a compact, orientable, hyperbolizable 3-manifold, let $\{\rho_n\} \subset \text{SH}(M)$ be a sequence of representations uniformizing M and let $\Gamma_n = (F_n, m_n, L_n)$ be the end invariants of ρ_n . Assume that $\{\Gamma_n\}$ converges to a doubly incompressible gallimaufry $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$. We proved in the preceding section that a subsequence of $\{\rho_n\}$ converges algebraically. We now show that this subsequence converges strongly to its algebraic limit $\rho_\infty(\pi_1(M))$, by showing that the convex cores of the $\rho_n(\pi_1(M))$ converge to the convex core of $\rho_\infty(\pi_1(M))$.

4.1 The ends of M_∞

We start with the geometrically finite ends of $\rho_\infty(\pi_1(M))$.

Lemma 4.1. *Let F be a connected component of the moderate surface F_∞ of Γ_∞ . Then there is a convex pleated surface $f_\infty : F \rightarrow M_\infty = \mathbb{H}^3/\rho_\infty(\pi_1(M))$ homotopic to the inclusion $F \subset M$.*

Proof. For n sufficiently large, F is a subsurface of F_n ; let H_n be the connected component of F_n containing F . We know that the restrictions of the m_n to F converge to a complete hyperbolic metric m on F . Let C_{ρ_n} be the convex core of $\rho_n(\pi_1(M))$. Since H_n is a component of F_n , there is a convex pleated surface $f_n : H_n \rightarrow \partial C_{\rho_n}$. Furthermore, by [EpM], H_n endowed with the metric induced by f_n is bilipschitz to (H_n, m_n) with a uniform bilipschitz constant. It follows that $\ell_{f_n}(\partial \bar{F}) \rightarrow 0$ and that $\{\ell_{f_n}(c)\}$ is bounded for any simple closed curve $c \subset F$, where ℓ_{f_n} is the length function of the metric induced by f_n . From this we deduce that there is a subsequence such that the restrictions to F of the metrics induced by the f_n converge to a complete hyperbolic metric.

Let $c \subset F$ be a simple closed curve such that $\rho_\infty(c)$ is a hyperbolic isometry. Since $\{\ell_{f_n}(c)\}$ is bounded, the distance between $c^* \subset M_n$ and $f_n(c)$ is uniformly bounded. Using Arzela-Ascoli's Theorem as in [CEG] we can extract a subsequence of $\{f_n\}$ that converges to a pleated surface $f_\infty : F \rightarrow C_{\rho_\infty}$ homotopic to the inclusion $F \subset M$. Since the f_n are convex surfaces, by [BoO], f_∞ is a convex surface as well, see also [Le2]. $\square$

Next we show that each component of the immoderate lamination L_∞ of Γ_∞ is an ending lamination of M_∞ .

Lemma 4.2. *Let L be a connected component of L_∞ . Then there is a geometrically infinite end E of $M_\infty = \mathbb{H}^3/\rho_\infty(\pi_1(M))$ such that E is homeomorphic to $S(L) \times [0, \infty)$, the inclusion $E \rightarrow M_\infty$ is homotopic to the inclusion $S(L) \hookrightarrow M$ and L is the ending lamination of E .*

Proof. Let S be the component of ∂M containing L and let $\sigma_n : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ be the representation induced from ρ_n by the inclusion map. We note that $\ell_{\sigma_n}(\mu^*) = \ell_{\rho_n}(\mu^*)$ for any measured geodesic lamination $\mu \in \mathcal{ML}(S) \subset \mathcal{ML}(\partial M)$.

By Claim 3.5, there is a sequence $\{c_n\}$ of simple closed curves converging in the Hausdorff topology to a geodesic lamination containing L such that $\{\ell_{\sigma_n}(c_n^*)\}$ is a bounded sequence. Choose a transverse measure λ supported by L . Since $\{\ell_{\sigma_n}(c_n^*)\}$ is a bounded sequence, it follows from the continuity of the length function, see [Bro], that $\ell_{\sigma_\infty}(\lambda^*) = 0$. This means that L is not realized in $\mathbb{H}^3/\sigma_\infty(\pi_1(S))$. It follows that L is an ending lamination of an end E' of $\mathbb{H}^3/\sigma_\infty(\pi_1(S))$.

By [Ca], the end E' covers an end E of M_∞ and the covering $E' \rightarrow E$ is finite to one. On the other hand, E' is homeomorphic to $S(L) \times [0, \infty)$ (this comes from the fact that the ending lamination has to "fill up" the surface defining the end) and if we consider the surface $S(L) \times \{1\}$ its image in E under the covering $E' \rightarrow E$ is homotopic to the inclusion $S(L) \hookrightarrow M$. Therefore the covering $E' \rightarrow E$ is a homeomorphism. Thus we have proved that there is a geometrically infinite end E of $M_\infty = \mathbb{H}^3/\rho_\infty(\pi_1(M))$ such that E is homeomorphic to $S(L) \times [0, \infty)$, the inclusion $E \hookrightarrow M_\infty$ is homotopic to the inclusion $S(L) \hookrightarrow M$ and L is the ending lamination of E . □

4.2 Reconstructing the convex core

In this section we show how the results of the preceding section allow us to describe the convex core of M_∞ .

Let L be a component of L_∞ . By Lemma 4.2, M_∞ has a geometrically infinite end homeomorphic to $S(L) \times [0, \infty)$ with ending lamination L . Choose some $p > 0$ and consider the map $f_\infty : S(L) \rightarrow S(L) \times \{p\}$. Each cusp of $S(L)$ is mapped under f_∞ to a cusp of M_∞ . Let G_∞ be the union of the surfaces $S(L)$ when L runs through all the components of L_∞ . We have thus constructed an embedding $f_\infty : G_\infty \rightarrow M_\infty$ which is homotopic to the inclusion map.

In Lemma 4.1, we defined a map $f_\infty : F_\infty \rightarrow C_{\rho_n}$ which is a homeomorphism onto its image and is homotopic to the inclusion map. Now we have a map $f_\infty : F_\infty \cup G_\infty \rightarrow M_\infty$. The complementary regions of $F_\infty \cup G_\infty$ in $\partial_{\chi < 0} M$ are annuli. By Lemmas 4.1 and 4.2, the simple closed curves in the homotopy class defined by each of these annuli corresponds to a parabolic element of $\rho_\infty(\pi_1(M))$. Furthermore, since Γ_∞ is doubly incompressible, to each such parabolic element of $\rho_\infty(\pi_1(M))$ there corresponds exactly one component of $\partial M - (F_\infty \cup G_\infty)$. To each such component (which is an annulus) corresponds two cusps of $F_\infty \cup G_\infty$ whose images under f_∞ are two homotopic non-compact annuli lying in a cusp of M_∞ . Remove from $f_\infty(F_\infty \cup G_\infty)$ these two non-compact annuli and connect the

boundary components so created by a compact annulus. Perform the same operation for all the components of $\partial_{\chi < 0} M - (F_\infty \cup G_\infty)$. We get a compact surface $S_\infty \subset M_\infty$.

Change f_∞ to get a homeomorphism $g_\infty : \partial_{\chi < 0} M \rightarrow S_\infty$ (this only involves making the correct choice of the Dehn twisting in $\partial_{\chi < 0} M - (F_\infty \cup G_\infty)$ so that g_∞ is homotopic to the inclusion $\partial_{\chi < 0} M \hookrightarrow M_\infty$). Adjoin to S_∞ the boundary of the rank 2 cusps of M_∞ and extend g_∞ to ∂M . Now we have a homeomorphism $g_\infty : \partial M \rightarrow S_\infty$ which is homotopic to the identity. Since ∂M bounds a compact 3-manifold, so does $g_\infty(\partial M)$. We deduce easily from this that $f_\infty(F_\infty \cup G_\infty)$ bounds a subset C_∞ of M_∞ which has a finite volume (compare with [BoO, Lemme 21]). Since $f_\infty(S(L_\infty))$ bounds an union of geometrically infinite ends and since $f_\infty(F_\infty)$ is an union of convex pleated surfaces, the union of C_∞ and of the geometrically infinite ends contains the convex core C_{ρ_∞} of ρ_∞ . If one component of ∂C_{ρ_∞} were to lie in $\text{int}(C_\infty)$, the corresponding geometrically finite end would lie inside C_∞ . This would contradict the fact that C_∞ has finite volume (see [BoO, Lemme 21] for more details). It follows that the union of C_∞ and of the geometrically infinite ends of M_∞ is the convex core of ρ_∞ .

Extend g_∞ to a homotopy equivalence h from M to the compact set bounded by $g_\infty(\partial M)$. By [Wa], h is homotopic to a homeomorphism. It follows that ρ_∞ uniformizes M .

4.3 Domain of discontinuity and strong convergence

Now that we know the geometrically finite ends and the behavior of the m_n , we can deduce from [Kl] that the algebraically convergent sequence $\{\rho_n\}$ converges strongly. On the other hand, as we have already proved the convergence of the convex cores, it is not hard now to conclude that we have strong convergence. So we briefly describe here why the sequence $\{\rho_n\}$ converges strongly.

The main tool is a result of [JoM] (see also [MaT]):

Theorem E. *Let $\{\rho_n\} \subset \text{AH}(M)$ be a sequence of representations that converges algebraically to ρ_∞ . Assume that Ω_{ρ_∞} is not empty. If $\{\Omega_{\rho_n}\}$ converges to Ω_{ρ_∞} in the sense of Carathéodory, then $\{\rho_n\}$ converges strongly to ρ_∞ .*

Recall that $\{\Omega_{\rho_n}\}$ converges to Ω_{ρ_∞} in the sense of Carathéodory if and only if $\{\Omega_{\rho_n}\}$ satisfies the two following conditions:

- every compact subset $K \subset \Omega_{\rho_\infty}$ lies in Ω_{ρ_n} for all sufficiently large n ;
- every open set O which lies in Ω_{ρ_n} for infinitely many n also lies in Ω_{ρ_∞} .

We show that the Ω_{ρ_n} and Ω_{ρ_∞} satisfy these two conditions.

Lemma 4.3. *Under the hypotheses of Proposition 3.1, a subsequence of $\{\rho_n\}$ converges strongly to ρ_∞ .*

Proof. When Ω_{ρ_∞} is empty, $\{\rho_n\}$ converges strongly to ρ_∞ by [Ca] (see [Kl]). Assume now that Ω_{ρ_∞} is not empty.

Consider the convex pleated surface $f_n : F_n \rightarrow C_{\rho_n}$. Let Π be a hyperbolic plane that

intersects $f_\infty(F_\infty)$ in a non-degenerate subsurface. The ideal boundary of Π in $\widehat{\mathbb{C}} = \partial\overline{\mathbb{H}}^3$ is a circle which bounds a disc $D \subset \partial\overline{\mathbb{H}}^3$ such that $\text{int}(D) \subset \Omega_{\rho_\infty}$.

Let $K \subset \Omega_{\rho_\infty}$ be a compact connected subset. Such a compact set is covered by the interiors of finitely many discs D_i defined as above. Since $\{f_n\}$ converges to f_∞ , each such disc D_i is the limit of a sequence $\{D_{i,n}\}$ where $D_{i,n} \subset \Omega_{\rho_n}$ is the disc bounded by the ideal boundary of a support plane for $f_n(F_n)$. It follows that for n sufficiently large, K is covered by the $D_{i,n}$. In particular, we have $K \subset \Omega_{\rho_n}$ for n sufficiently large.

Now, we will prove, by contradiction, that every open set O which lies in Ω_{ρ_n} for infinitely many n also lies in Ω_{ρ_∞} . Let O be an open set lying in Ω_{ρ_n} for infinitely many n and let $w \in O \cap \Lambda_{\rho_\infty}$. Since the fixed points of hyperbolic isometries are dense in Λ_{ρ_∞} and since $\{\rho_n\}$ converges to ρ_∞ , there is a sequence $\{w_n\}$ of points converging to w such that $w_n \in \Lambda_{\rho_n}$. For n sufficiently large, we have $w_n \in O \cap \Lambda_{\rho_n}$, contradicting our hypothesis that $O \subset \Omega_{\rho_n}$ for infinitely many n . $\square$

Finally, we have now proved that the ending map is proper, namely:

Proposition 4.4. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\{\rho_n\} \subset \text{AH}(M)$ be a sequence of representations uniformizing M and let $\Gamma_n = (F_n, m_n, L_n)$ be the end invariants of ρ_n . Assume that $\{\Gamma_n\}$ converges in $\mathcal{GA}(\partial M)$ to a doubly incompressible gallimaufry $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$. Then a subsequence of $\{\rho_n\}$ converges strongly.* $\square$

5 Necessary conditions

In this section we prove that the end invariants of a sequence $\{\rho_n\}$ converge to the end invariants of the limit.

Proposition 5.1. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\{\rho_n\}$ be a sequence of representations that uniformize M . Assume that ρ_n is geometrically finite and minimally parabolic for all n and let $\Gamma_n = (\partial_{\chi < 0} M, m_n, \emptyset)$ be its end invariants. If $\{\rho_n\}$ converges strongly to a representation ρ_∞ , then $\{\Gamma_n\}$ converges in $\mathcal{GA}(\partial M)$ to the end invariants Γ_∞ of ρ_∞ .*

Proof. First recall that we have seen in Lemma 2.1 that ρ_∞ uniformizes M . Let us also recall the topology of $\mathcal{GA}(\partial M)$ as described in Section 2.8. Consider our sequence of gallimaufries $\{\Gamma_n = (\partial_{\chi < 0} M, m_n, \emptyset)\}$. Then $\{\Gamma_n\}$ converges to $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$ in $\mathcal{GA}(\partial M)$ if we have the following:

- i) for any n we have $F_\infty \subset F_n$ and the restrictions of the m_n to F_∞ converge to m_∞ , namely for any closed curve $c \subset F_\infty$, we have $\ell_{m_n}(c) \rightarrow \ell_{m_\infty}(c)$,
- iii) if a component L of L_∞ lies in infinitely many of the F_n , then the restrictions of the m_n to $S(L)$ tend to L .

Notice that since the Γ_n lie in $\text{int}(\mathcal{GA}(\partial M))$ the definitions are simpler than in the general case. In particular, part (ii) of the definition is trivially satisfied.

The first property can be deduced from the convergence of the limit sets.

Lemma 5.2. *Consider the moderate surface F_∞ of Γ_∞ equipped with its complete hyperbolic metric m_∞ . For any closed curve $c \subset F_\infty$, we have $\ell_{m_n}(c) \longrightarrow \ell_{m_\infty}(c)$.*

Proof. Let F be the component of F_∞ containing c and let S be the component of ∂M containing F . Since $\{\rho_n\}$ converges strongly, as proved in [KeT], the limit sets $\{\Lambda_{\rho_n}\}$ of the Kleinian groups $\rho_n(\pi_1(S))$ converge to the limit set of $\rho_\infty(\pi_1(S))$ in the Hausdorff topology. It follows that $\{\Omega_{\rho_n}\}$ converges to Ω_{ρ_∞} in the sense of Carathéodory, and hence that the Poincaré metrics on the Ω_{ρ_n} converge to the Poincaré metric on Ω_{ρ_∞} . By this, we mean that if we consider a point $x \in \Omega_{\rho_\infty}$, and a sequence $\{x_n\} \subset \Omega_{\rho_n}$ converging to x (whose existence is guaranteed by the convergence of the Λ_{ρ_n}) the Poincaré metric on Ω_{ρ_n} at x_n converges to the Poincaré metric on Ω_{ρ_∞} at x , see [He].

Consider a component O of Ω_{ρ_∞} covering F and an arc $\tilde{c}$ that is a lift of c to O . Let $G_O \subset \pi_1(M)$ be the subgroup such that $\rho_\infty(G_O)$ is the stabilizer of O . If we let x be one endpoint of $\tilde{c}$, then the other endpoint is the point $\rho_\infty(g)(x)$, where $\rho_\infty(g)$ is the element of $\rho_\infty(G_O)$ (up to conjugacy) corresponding to c . Let K be the closure of an ε -neighborhood of $\tilde{c}$, and note that K is a compact subset of O . In particular, $K \subset \Omega_{\rho_n}$ for all n sufficiently large. Since the Poincaré metrics on the Ω_{ρ_n} converge to the Poincaré metric on Ω_{ρ_∞} uniformly on compact subsets, since $\rho_n(g)(x)$ lies in K for n sufficiently large and since $\{\rho_n(g)(x)\}$ converges to $\rho_\infty(g)(x)$, we see that $\ell_{m_n}(c) \longrightarrow \ell_{m_\infty}(c)$, as desired. $\square$

Next we prove that for a connected component L of L_∞ , up to extracting a subsequence, the restriction of m_n to $S(L)$ tends to a measured geodesic lamination supported by L . To prove this property, we use some of the ideas of the "lemme d'intersection" [Bon, Proposition 3.4]. Consider a component L of the immoderate lamination of Γ_∞ . By definition of the end invariants of ρ_∞ , L is the ending lamination of a geometrically infinite end of M_∞ . We want to show that, up to extracting a subsequence, the restrictions of the m_n to $S(L)$ tend to a measured geodesic lamination supported by L .

Lemma 5.3. *Consider a connected component L of L_∞ . Then, up to extracting a subsequence, the restriction of the m_n to $S(L)$ tend to a measured geodesic lamination supported by L .*

Before starting the proof of Lemma 5.3 recall that it finishes the proof of Proposition 5.1. Putting Lemmas 5.2 and 5.3 together, we conclude that given a strongly convergent sequence $\{\rho_n\}$ of representations that uniformize M with end invariants $\Gamma_n = (\partial_{\chi < 0} M, m_n, \emptyset)$, then $\{\Gamma_n\}$ converges in $\mathcal{GA}(\partial M)$ to the end invariants of the limit ρ_∞ . This concludes the proof of Proposition 5.1. $\square$

Proof of Lemma 5.3. Let S be the connected component of ∂M that contains L . Since L is an irrational minimal component of the immoderate part of L_∞ it is the ending lamination of a geometrically infinite end of M_∞ . It follows, see [Th1] and [Bon], that there is a sequence $\{e_k\} \subset \partial M$ of simple closed curves such that $\{e_k\}$ converges in $\mathcal{PML}(\partial M)$ to a projective

measured geodesic lamination supported by L , the geodesic representatives $e_{k,\infty}^*$ of the e_k in M_∞ exit any compact subset of M_∞ , and $\{\ell_{\rho_\infty}(e_{k,\infty}^*)\}$ is bounded.

Since $\{\rho_n\}$ converges strongly to ρ_∞ , there are sequences $q_n \rightarrow 1$ and $R_n \rightarrow \infty$, a point $x_\infty \in M_\infty$, and a sequence of diffeomorphisms $\psi_n : B(x_\infty, R_n) \subset M_\infty \rightarrow M_n$ such that $\psi_n(x_\infty) = x_n$ and that ψ_n is a q_n -bilipschitz diffeomorphism onto its image. Choose ε such that for any k either $e_{k,\infty}^*$ is disjoint from the ε -thin part of M_∞ or $e_{k,\infty}^*$ is the core of a Margulis tube.

Consider a sequence $\{P_n\}$ of pants decompositions of S such that $\{\ell_{m_n}(P_n)\}$ is a bounded sequence. We show that, up to extracting a subsequence, $\{P_n\}$ converges in the Hausdorff topology to a geodesic lamination containing L . Then we show that this can happen only when the m_n degenerate transversely to L . First we control the intersection numbers between P_n and e_k using the following Lemma.

Lemma 5.4. *Consider a sequence $\{d_n\} \subset S$ of simple closed curves. There is $Q > 0$ such that for any k sufficiently large there is N_k such that $i(e_k, d_n) \leq Q(\ell_{m_n}(d_n) + 2\pi)$ for $n \geq N_k$.*

Proof. The basis of this proof comes from [Bon, Proposition 3.4].

Let $K_\infty \subset C_{\rho_\infty} \subset M_\infty = \mathbb{H}^3/\rho_\infty(\pi_1(M))$ be a compact core for M_∞ . Since $\{\rho_n\}$ converges strongly to ρ_∞ , for n sufficiently large $\psi_n(K_\infty)$ is a compact core for M_n (see Lemma 2.1). By [KeT], $\{\Lambda_{\rho_n}\}$ converges to Λ_{ρ_∞} in the Hausdorff topology. By [Bow], the convex hulls $\{H_{\rho_n}\} \subset \mathbb{H}^3$ converge to $H_{\rho_\infty} \subset \mathbb{H}^3$ in the Hausdorff topology. It follows that we have $\psi_n(K_\infty) \subset C_{\rho_n}$ for n large enough. We use the notation $K_n = \psi_n(K_\infty)$. Notice that the induced metrics on the ∂K_n converge to the induced metric on ∂K_∞ .

Recall that $e_{k,n}^*$ is the geodesic representative of e_k in M_n . Since $e_{k,\infty}^*$ exits any compact set, $d(e_{k,\infty}^*, K_\infty) \geq 3C_k$ for some $C_k \rightarrow \infty$. Fix k . For n sufficiently large, $e_{k,\infty}^*$ lies in $B(x_\infty, R_n) \subset M_\infty$. Furthermore $\ell_{\rho_n}(\psi_n(e_{k,\infty}^*)) \leq q_n \ell_{\rho_\infty}(e_{k,\infty}^*)$ where q_n is the bilipschitz constant of ψ_n . So $\ell(e_{k,n}^*) \leq q_n \ell_{\rho_\infty}(e_{k,\infty}^*)$ for n sufficiently large. Since the length of $e_{k,\infty}^*$ is bounded, there are Q_1 and N_k such that for $n \geq N_k$, we have $\ell_{\rho_n}(e_{k,n}^*) \leq Q_1$ (note that Q_1 depends on the sequence e_k but not on k). For n sufficiently large, $d(\psi_n(e_{k,\infty}^*), K_n) \geq 2C_k$. Since ψ_n is q_n -bilipschitz, $\psi_n(e_{k,\infty}^*)$ is a q_n^2 -quasi-geodesic. It follows that $d(e_{k,n}^*, K_n) \geq C_k$ for n sufficiently large.

In what follows, we view d_n and e_k as simple closed curves in ∂K_∞ . Let $f_n : S \rightarrow \partial C_{\rho_n}$ be the map sending $S \subset \partial M$ to the corresponding part of the boundary of the convex core of $M_n = \mathbb{H}^3/\rho_n(\pi_1(M))$. Consider an annulus A_{d_n} joining $\psi_n(d_n)$ to $f_n(d_n)$ in C_{ρ_n} and an annulus $A_{e_{k,n}}$ joining $\psi_n(e_k)$ to $e_{k,n}^*$ in $\text{int}(C_{\rho_n})$. By [Bon, Lemme 3.2], we have $i(e_k, d_n) \leq i(f_n(d_n), A_{e_{k,n}}) + i(e_{k,n}^*, A_{d_n})$ where the first term is the geometric intersection number in S and the two others are geometric intersection numbers in M_n . Since $A_{e_{k,n}}$ lies in $\text{int}(C_{\rho_n})$ it does not intersect ∂C_{ρ_n} . It follows then that $i(f_n(d_n), A_{e_{k,n}}) = 0$. Hence we only need to bound $i(e_k, A_{d_n})$.

For each n , we choose a point $x_n \subset A_{d_n}$ such that $\{d(K_n, x_n)\}$ is a bounded sequence. We also choose points $y_n \subset \psi_n(d_n)$ and $z_n \subset f_n(d_n)$. Consider an annulus $A_{d_n} \subset C_{\rho_n}$ joining $\psi_n(d_n)$ to $f_n(d_n)$. Let us change A_{d_n} by a homotopy so that it contains x_n and a geodesic segment k_n joining y_n to z_n . Cutting A_{d_n} along k_n , we get a disc D_n containing x_n . Let us change D_n by a homotopy to the geodesic from x_n to ∂D_n . Gluing back D_n along k_n we

get an annulus A_{d_n} joining $\psi_n(d_n)$ to $f_n(d_n)$. This annulus A_{d_n} is the union of the set E_n of geodesic segments joining $f\psi_n(d_n)$ to x_n , of the set F_n of geodesic segments joining k_n to x_n and of the set G_n of geodesic segments joining $f_n(d_n)$ to x_n . According to the previous paragraph, we have $i(e_k, d_n) \leq i(e_{k,n}^*, A_{d_n})$.

By the choice of x_n , for n sufficiently large, $E_n \subset \psi_n(B(x_\infty, r_n))$ and $\psi_n^{-1}(E_n)$ lies in a compact set (independent of n). It follows that for k sufficiently large, $e_{k,n}^*$ does not intersect E_n . Thus for k sufficiently large, $e_{k,n}^*$ only intersects $H_n = F_n \cup G_n$. Since $d(\psi - n^{-1}(E_n), e_{k,\infty}^*) \xrightarrow{k \rightarrow \infty} \infty$, we even have some $\varepsilon > 0$ such that for n sufficiently large the intersections between $e_{k,n}^*$ and H_n are at distance at least ε from the two geodesic segments in H_n joining x_n to y_n . Furthermore, since $\{H_{\rho_n}\}$ converges to H_{ρ_∞} in the Hausdorff topology, we have $d(e_{k,n}^*, \partial C_{\rho_n}) \geq \varepsilon$, for n large enough. Thus the intersections between $e_{k,n}^*$ and H_n are at distance at least ε from ∂H_n .

By construction, F_n is the union of two geodesic triangles. Thus we have $\text{area}(H_n) \leq 2\pi$. Since $f_n(d_n)$ is piecewise geodesic, G_n is an union of geodesic triangles. It is well known that the area of a hyperbolic triangle is less than the length of any of its edges, see [Th1, Lemma 9.3.2]. Thus we have $\text{area}(G_n) \leq \ell(f_n(d_n))$.

From the proof of [Bon, Proposition 3.4], we get:

- If $\ell(e_{k,n}^*) > \varepsilon$, then $i(e_k, d_n) \leq Q_2^{-1} \ell_{\rho_n}(e_{k,n}^*) \text{area}(H_n)$, where Q_2 is the volume of the ball with radius $\frac{\varepsilon}{8}$ in $\mathbb{R} \times \mathbb{H}^2$.

- If $\ell(e_{k,n}^*) \leq \varepsilon$, then we have $i(e_k, d_n) \leq Q_3^{-1} \text{area}(H_n)$, where Q_3 is the area of a hyperbolic disc with radius $\frac{\varepsilon}{8}$.

We have seen above the inequality

$$\text{area}(H_n) \leq \ell(f_n(d_n)) + 2\pi$$

By [Ah], we have $\ell(f_n(d_n)) \leq 2\ell_{m_n}(d_n)$. Consider $Q = \max\{2Q_1Q_2^{-1}, 2Q_3^{-1}\}$ and notice that Q depends on the choice of the sequence e_k . For k sufficiently large and for n sufficiently large (depending on k), we have $i(e_k, d_n) \leq Q(\ell_{m_n}(d_n) + 2\pi)$. $\square$

We can now show that any sequence of short pants decompositions, with respect to the m_n , converges to L . More precisely, we have:

Claim 5.5. *Let $\{P_n\}$ be a sequence of pants decompositions of S such that $\{\ell_{m_n}(P_n)\}$ is a bounded sequence. Extract a subsequence such that $\{P_n\}$ converges in the Hausdorff topology to a geodesic lamination $P_\infty \subset S$. Then L is a sublamination of P_∞ .*

Proof. Since L is an irrational geodesic lamination, if P_∞ crosses L , then we have $i(e_k, P_n) \rightarrow \infty$. By Lemma 5.4, $i(e_k, P_n)$ is bounded. It follows that either L is disjoint from P_∞ or L is a sublamination of P_∞ . Since P_n is a pants decomposition for any n , no geodesic lamination is disjoint from P_∞ . So we conclude that L is a sublamination of P_∞ . $\square$

Now we can conclude the proof of Lemma 5.3. As we have seen, there is a constant K depending only on S such that for any complete hyperbolic metric m on S , there is a

pants decomposition P_S with $\ell_m(P_S) \leq K$ (see [Be]). Consider a sequence $\{P_n\}$ of pants decompositions such that $\{\ell_{m_n}(P_n)\}$ is bounded. Extract a subsequence such that $\{P_n\}$ converges in the Hausdorff topology to a geodesic lamination P_∞ . By Claim 5.5, L is a sublamination of P_∞ . Let $c \subset S(L)$ be a non-peripheral simple closed curve. Since c crosses L , we have $i(P_n, c) \rightarrow \infty$. It follows then from the Collar Lemma that $\ell_{m_n}(c) \rightarrow \infty$. In particular the restriction of m_n to $S(L)$ is unbounded. By Lemma 3.4, the restriction of m_n to $S(L)$ tends to L . This concludes the proof of Lemma 5.3 and of Proposition 5.1. $\square$

Using the Ending Lamination Classification, Propositions 4.4 and 5.1, we can now prove Theorem D.

Theorem D. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. The ending map that to a representation uniformizing M associates its end invariants is a homeomorphism into the set of doubly incompressible gallimaufries.*

Proof. By the Ending Lamination Classification, see [Mi2] and [BCM], the ending map is injective. By Proposition 4.4, this map is proper. Combining Propositions 4.4 and 5.1, we get that a sequence $\{\rho_n\} \subset \text{SH}(M)$ with end invariants $\{\Gamma_n = (\partial_{\chi < 0} M, m_n, \emptyset)\}$ converges strongly to ρ_∞ with end invariants $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$ if and only if $\{\Gamma_n\}$ converges to Γ_∞ in $\mathcal{GA}(\partial M)$. In particular it follows that any $\rho_\infty \in \text{SH}(M)$ can be approximated by geometrically finite minimally parabolic representations in the strong topology, namely $\text{SH}(M)$ is the closure of its interior. By using such approximations we can drop the assumption in Proposition 5.1 that ρ_n is geometrically finite and minimally parabolic. It follows that the ending map is continuous. Thus the ending map is injective, continuous and proper: it is a homeomorphism onto its image. $\square$

From this Theorem and from Lemma 2.4, we deduce that $\text{SH}(\pi_1(M))$ is not locally compact (Lemma 1.1).

Lemma 1.1. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Then the space $\text{SH}(M)$ is not locally compact.* $\square$

6 Self bumping and local connectivity

By constructing appropriate paths in $\mathcal{GA}(\partial M)$ we study the local connectivity of the set of doubly incompressible gallimaufries. We note that by [Le1], the set of doubly incompressible gallimaufries is an open subset of $\mathcal{GA}(\partial M)$.

A note about the notation in this section: The approximating gallimaufries all have the same moderate surface, namely $\partial_{\chi < 0}(\partial M)$, and the same immoderate lamination, namely the empty lamination. Therefore, we identify each approximating gallimaufry with the metric on $\partial_{\chi < 0}(\partial M)$. In particular, we speak of sequences of metrics on the moderate surface as converging to a limiting gallimaufry.

Proposition 6.1. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\Gamma = (F, m, L)$ be a doubly incompressible gallimaufry and let $\{(\partial_{\chi < 0} M, m_n, \emptyset)\}$ and $\{(\partial_{\chi < 0} M, s_n, \emptyset)\}$ be two sequences $\text{int}(\mathcal{GA}(\partial M))$ converging to Γ . Then, up to passing to*

a subsequence, there is an arc $k_n \subset \text{int}(\mathcal{GA}(\partial M))$ joining m_n to s_n such that any sequence of points $\{z_n \in k_n\}$ converges to Γ .

Proof. The main difficulty is that on F we need to have precise control on the behavior of z_n , whereas on $S(L)$ we need to control the large scale behavior of z_n . To deal with that issue, we use Fenchel-Nielsen coordinates to control the behavior of z_n on F and the measured geodesic lamination $\mathcal{F}_P(z_n)$ as defined in section 2.7 to control the large scale behavior of z_n on $S(L)$.

We denote by G the surface obtained by adding to $\overline{F}$ any component of $\partial M - \overline{F}$ that is an essential annulus. For each n and t we will define the metric $k_n(t)$ independently on G and on the closure H of $\partial_{\chi < 0} M - G$.

Choose $\varepsilon > 0$ and consider a pants decomposition P of ∂M so that any component of $\partial \overline{F}$ is homotopic to a leaf of P and that we have $\ell_{m_n}(c) \geq \varepsilon$ and $\ell_{s_n}(c) \geq \varepsilon$ for any n and any component c of P that is non-peripheral in $\overline{F}$.

Let us first define the restriction of $k_n(t)$ to H . In $(\partial_{\chi < 0} M, m_n, \emptyset)$ and $(\partial_{\chi < 0} M, s_n, \emptyset)$ we view H as a subsurface with geodesic boundary. Since s_n and m_n tend to Γ , then we have $\ell_{m_n}(\partial H) \rightarrow 0$ and $\ell_{s_n}(\partial H) \rightarrow 0$. Then for n large enough, there is associated to the restriction of m_n , respectively s_n , to H a measured geodesic lamination $\mathcal{F}_P(m_n)$, respectively $\mathcal{F}_P(s_n)$ (see section 2.7). Furthermore the restriction of m_n to H is uniquely defined by $\mathcal{F}_P(m_n)$ and $\{\ell_{m_n}(c) | c \text{ is a component of } \partial H\}$.

Extract a subsequence such that $\{\mathcal{F}_P(m_n)\}$ converges to a geodesic lamination R in the Hausdorff topology. It follows from Theorem 2.2 that L is a sublamination of R . Consider a sequence $\{\tau_n\}$ of train tracks carrying R so that $\{\tau_n\}$ is a basis of neighborhood of R for the Hausdorff topology. Namely any sequence of laminations $\{R_n\}$ such that R_n is minimally carried by τ_n converges to R in the Hausdorff topology (for the existence of such train tracks, see [Ot1]). Notice that if R_n is only carried by τ_n then up to extracting a subsequence $\{R_n\}$ converges to a sublamination of R . Since L is a sublamination of R , we may choose the τ_n so that L is minimally carried by a sub-track of τ_n for any n .

Since $\{\mathcal{F}_P(m_n)\}$ converges to R , up to changing the indices, we may assume that $\mathcal{F}_P(m_n)$ is carried by τ_n . In particular $\mathcal{F}_P(m_n)$ defines a weight system $\mathcal{F}_P(m_n)(\tau_n)$ on τ_n . Let λ be a measured geodesic lamination with support L , and note that it defines a weight system $\lambda(\tau_n)$. Let $\{K_n\}$ be a sequence of positive real number tending to ∞ . For $t \in [0, \frac{1}{2}]$, the weight system $2(\frac{1}{2} - t)\mathcal{F}_P(m_n)(\tau_n) + 2tK_n\lambda(\tau_n)$ defines a measured geodesic lamination $\mu_n(t)$ carried by τ_n such that $\mu_n(0) = \mathcal{F}_P(m_n)$ and $\mu_n(\frac{1}{2}) = K_n\lambda$. For $t \neq \frac{1}{2}$, μ_n is minimally carried by τ_n . It follows that we have:

Claim 6.2. *Consider a sequence $\{t_n\} \in [0, \frac{1}{2}]^{\mathbb{N}}$. Then the Hausdorff limit of any subsequence of $\{\mu_n(t_n)\}$ is either R or L . In particular it contains L .*

Consider a simple closed curve $c \subset G$. The intersection number $i(c, \mu_n(t))$ is given by the following formula:

Claim 6.3. *For n large enough, depending on c , we have $i(c, \mu_n(t)) = 2ti(c, \mathcal{F}_P(m_n)) + 2(\frac{1}{2} - t)K_n i(c, \lambda_n)$ for any $t \in [0, 1]$.*

Proof. Since $\{\tau_n\}$ is a basis of neighborhoods for R for the Hausdorff topology, for n large enough, depending on c , we can change τ_n and c by homotopies so that for any measured geodesic lamination λ carried by τ_n , $i(c, \lambda)$ is the sum of the weights of the branches of τ_n that c intersects transversely, counted with multiplicity. In particular, for n large enough, we have $i(c, \mu_n(t)) = 2(\frac{1}{2} - t)i(c, \mathcal{F}_P(m_n)) + 2tK_n i(c, \lambda_n)$. $\square$

Using a similar construction, replacing m_n by s_n we get a path of measured geodesic laminations $\mu_n(t)$, $t \in [\frac{1}{2}, 1]$ with $\mu_n(\frac{1}{2}) = K_n \lambda_n$ and $\mu_n(1) = \mathcal{F}_P(s_n)$. Hence, given a sequence $\{t_n\} \in [\frac{1}{2}, 1]^n$, the Hausdorff limit of any subsequence of $\{\mu_n(t_n)\}$ contains L .

We define the restriction of $k_n(t)$ to H for $t \in [0, 1]$ as follows : $\ell_{k_n(t)}(c) = (1-t)\ell_{m_n}(c) + t\ell_{s_n}(c)$ for any component c of ∂H and $\mathcal{F}_P(k_n(t)) = \mu_n(t)$.

Next we define the restriction of $k_n(t)$ to G . Consider the Fenchel-Nielsen coordinates $\{(\ell_c, \theta_c) | c \text{ a leaf of } P\}$ associated to P . Namely for a metric m , $\ell_c(m) \in]0, \infty[$ is the m -length of c and $\theta_c \in \mathbb{R}$ is the twist angle along c (taking the convention that $\theta_c(m') = \theta_c(m) + 2\pi$ corresponds to a full Dehn twist along c). On G we define the restriction of $k_n(t)$ as follows : each component c of ∂G is a closed geodesic with length $\ell_c(k_n(t)) = (1-t)\ell_c(m_n) + t\ell_c(s_n)$; the Fenchel-Nielsen coordinates of $k_n(t)$ along each non-peripheral leaf c of $P \cap G$ are $\{(1-t)\ell_c(m_n) + t\ell_c(s_n), (1-t)\theta_c(m_n) + t\theta_c(s_n)\}$. This defines a metric with geodesic boundary on G .

Notice that a component d of ∂G is also a component of ∂H and that the length given to d by the restriction of $k_n(t)$ to G is the same as the length given by the restriction to H . We just need to define how we glue together the metrics defined on G and H to get a complete definition of $k_n(t)$. For a component c of ∂G , the Fenchel-Nielsen coordinates of $k_n(t)$ along c are $\{(1-t)\ell_c(m_n) + t\ell_c(s_n), (1-t)\theta_c(m_n) + t\theta_c(s_n)\}$.

Now a complete hyperbolic metric $k_n(t)$ has been defined on $\partial_{X<0}M$ for any $n \in \mathbb{N}$ and any $t \in [0, 1]$. It follows easily from the definition that, for a fixed n , $k_n(t)$ is a continuous path joining m_n to s_n . The following two Claims will conclude the proof of Proposition 6.1.

Claim 6.4. *For any sequence $\{t_n\} \subset [0, 1]^{\mathbb{N}}$ the restriction of $k_n(t_n)$ to G tends to the complete hyperbolic metric m on F .*

Proof. Since $\{m_n\}$ and $\{s_n\}$ both converge to m , for any component c of P that is homotopic to a cusp of F , we have $\ell_{m_n}(c) \rightarrow 0$ and $\ell_{s_n}(c) \rightarrow 0$. Thus we get $\ell_{k_n(t_n)}(c) = t_n \ell_c(m_n) + (1-t_n)\ell_c(s_n) \rightarrow 0$. If c is a component of $P \cap G$ that is not homotopic to a cusp of F , then $\{\ell_c(m_n), \theta_c(m_n)\}$ and $\{\ell_c(s_n), \theta_c(s_n)\}$ converge to $\{\ell_c(m), \theta_c(m)\}$. Hence $\{\ell_c(k_n(t_n)), \theta_c(k_n(t_n))\} = \{t_n \ell_c(m_n) + (1-t_n)\ell_c(s_n), t_n \theta_c(m_n) + (1-t_n)\theta_c(s_n)\}$ converges to $\{\ell_c(m), \theta_c(m)\}$. It follows that the restriction of $k_n(t)$ to G tends to the complete hyperbolic metric m on F . $\square$

Claim 6.5. *Let L_1 be a component of L . Then for any sequence $\{t_n\} \in [0, 1]^{\mathbb{N}}$, the restriction of $k_n(t_n)$ to $S(L)$ tends to L .*

Proof. We will assume that we have $\{t_n\} \in [0, \frac{1}{2}]^{\mathbb{N}}$. The proof is similar if we assume $\{t_n\} \in [\frac{1}{2}, 1]^{\mathbb{N}}$ and these two results are enough to conclude the proof of Proposition 6.1.

Consider a non-peripheral simple closed curve $c \subset S(L_1)$. Since the restriction of m_n to $S(L_1)$ tends to L_1 , we have $i(\mathcal{F}_P(m_n), c) \rightarrow \infty$ and $\frac{i(\mathcal{F}_P(m_n), \partial S(L_1))}{i(\mathcal{F}_P(m_n), c)} \rightarrow 0$. We also have that $K_n i(\lambda_n, c) \rightarrow \infty$ and $K_n i(\lambda_n, d) = 0$. From Claim 6.3 and Theorem 2.2, we get $l_{k_n(t_n)}(c) \rightarrow \infty$ and $\frac{l_{k_n(t_n)}(\partial S(L_1))}{l_{k_n(t_n)}(c)} \rightarrow \infty$. Hence, up to extracting a subsequence the restriction of $k_n(t_n)$ to $S(L_1)$ tends to a projective measured geodesic lamination. By Claim 6.2, the support of this projective lamination is L_1 . This concludes the proof of Claim 6.4. $\square$

By Claims 6.4 and 6.5, for any sequence $\{t_n\} \in [0, 1]^{\mathbb{N}}$, $\{k_n(t_n)\}$ converges to Γ . $\square$

Using Theorem D and Proposition 6.1, we can now prove Theorems A and B.

Theorem A. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Then the space $\text{SH}(\pi_1(M))$ is locally connected.*

Proof. Consider a representation $\rho \in \text{SH}(\pi_1(M))$. By the Core theorem, see [Sc], there is a compact manifold M' such that ρ uniformizes M' . By Lemma 2.1, any sufficiently small neighborhood $\mathcal{V} \subset \text{SH}(\pi_1(M))$ of ρ lies in $\text{SH}(M')$. Let $\Gamma \in \mathcal{GA}(\partial M')$ be the end invariants of ρ . By Theorem D, $\mathcal{V}$ is homeomorphic to a neighborhood $\mathcal{W} \subset \mathcal{GA}(\partial M')$ of Γ . By Proposition 6.1, $\mathcal{W}$ contains a connected neighborhood $\mathcal{W}'$ of Γ . Taking the preimage of $\mathcal{W}'$ under the ending map, we get a connected neighborhood $\mathcal{V}' \subset \mathcal{V}$ of ρ . Thus we have proved that $\text{SH}(\pi_1(M))$ is locally connected. $\square$

The proof of Theorem B follows the same lines.

Theorem B. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $\rho \in \text{SH}(\pi_1(M))$ be a representation uniformizing M . Then any neighborhood of ρ contains a neighborhood $\mathcal{V} \subset \text{SH}(\pi_1(M))$ of ρ such that $\mathcal{V} \cap \text{int}(\text{SH}(\pi_1(M)))$ is connected.*

Proof. Let $\mathcal{V}_1 \subset \text{SH}(\pi_1(M))$ be a neighborhood of ρ . By Lemma 5.1, $\mathcal{V}_1$ contains a neighborhood $\mathcal{V}_2 \subset \text{SH}(M)$ of ρ . We denote by e the ending map as defined in Theorem D. By Proposition 6.1, $e(\mathcal{V}_2)$ contains a neighborhood $\mathcal{W} \subset \text{Im}(e)$ of $e(\rho)$ such that $\mathcal{W} \cap \text{int}(\mathcal{GA}(\partial M))$ is connected. Taking $\mathcal{V} = e^{-1}(\mathcal{W})$, we get a neighborhood of ρ such that $\mathcal{V} \cap \text{int}(\text{SH}(\pi_1(M)))$ is connected. $\square$

7 The action of $\text{Mod}(M)$

In this last section we study the action of $\text{Mod}(M)$ on $\text{SH}(M)$.

Theorem C. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Assume that M is not an I -bundle over a closed surface. Then the action of $\text{Mod}(M)$ on $\text{SH}(M)$ is properly discontinuous.*

Proof. Consider a sequence of representations $\{\rho_n\} \subset \text{int}(\text{SH}(M))$ and a sequence of diffeomorphisms $\{\psi_n : M \rightarrow M\}$ such that $\{\rho_n\}$ and $\{\rho_n \circ \psi_{n*}\}$ converge respectively to representations ρ_∞ and ρ'_∞ (up to conjugacy) in $\text{SH}(M)$. We show that up to extracting a subsequence, the ϕ_n are isotopic. It follows easily that the action of $\text{Mod}(M)$ on $\text{SH}(M)$ is properly discontinuous. Since minimally parabolic geometrically finite representations are dense in $\text{SH}(M)$, we may assume that ρ_n is such a representation without any loss of generality. As a first step, the following Claim is useful to pick the correct representation in each conjugacy class.

Claim 7.1. *Let $\{\gamma_n\}$, $\{a_n\}$ and $\{b_n\}$ be sequences of elements of $\text{PSL}(2, \mathbb{C})$. Assume that the sequences $\{a_n\}$ and $\{b_n\}$ converge to hyperbolic isometries a_∞ and b_∞ such that a_∞ and b_∞ do not have a common fixed point. Assume also that the sequences $\{\gamma_n a_n \gamma_n^{-1}\}$ and $\{\gamma_n b_n \gamma_n^{-1}\}$ converge. Then, up to extracting a subsequence, $\{\gamma_n\}$ converges.*

Proof. Assume that no subsequence of $\{\gamma_n\}$ converges. We show that we end up with a contradiction. For n sufficiently large, a_n and b_n are hyperbolic isometries. Let A_n and B_n be the axes of a_n and b_n respectively, for $n \in \mathbb{N} \cup \{\infty\}$. By assumption $\{A_n\}$, respectively $\{B_n\}$, converges to A_∞ , respectively B_∞ , in the Hausdorff topology on $\mathbb{H}^3 \cup \partial_\infty(\mathbb{H}^3)$. Consider a point $x \in \mathbb{H}^3$. Since we have assumed that no subsequence of $\{\gamma_n\}$ converges, we have $d(x, \gamma_n^{-1}(x)) \rightarrow \infty$. Since a_∞ and b_∞ do not have a common fixed point, we have $\max\{d(\gamma_n^{-1}(x), A_n), d(\gamma_n^{-1}(x), B_n)\} \rightarrow \infty$.

By assumption the sequence $\{d(x, \gamma_n a_n \gamma_n^{-1}(x))\}$ converges. On the other hand, we have $d(x, \gamma_n a_n \gamma_n^{-1}(x)) = d(\gamma_n^{-1}(x), a_n \gamma_n^{-1}(x))$ which converges if and only if $\{d(\gamma_n^{-1}(x), A_n)\}$ is a bounded sequence. Similarly we get that $\{d(\gamma_n^{-1}(x), B_n)\}$ is a bounded sequence. Thus we get that $\max\{d(\gamma_n^{-1}(x), A_n), d(\gamma_n^{-1}(x), B_n)\}$ is bounded, contradicting the preceding paragraph. $\square$

This Claim allows us to get the expected conclusion under some extra assumptions on the diffeomorphisms ϕ_n .

Lemma 7.2. *Let M be a compact, orientable, hyperbolizable 3-manifold with incompressible boundary. Let $N \subset \partial M$ be an incompressible compact submanifold such that $\pi_1(N)$ is not Abelian. Let $\{\phi_n : M \rightarrow M\}$ be a sequence of diffeomorphisms such that $\phi_n|_N$ is isotopic to the identity and let $\{\rho_n\} \subset \text{int}(\text{SH}(M))$ be a convergent sequence. Assume that the sequence $\{\rho_n \circ \phi_{n*}\}$ converges as well. Then there is a diffeomorphism $\phi : M \rightarrow M$ such that up to extracting a subsequence, each $\phi^{-1} \circ \phi_n$ is isotopic to the identity.*

Proof. Let $\rho_n : \pi_1(M) \rightarrow \text{PSL}(2, \mathbb{C})$ be a representative of $\rho_n \in \text{int}(\text{SH}(M))$ such that $\{\rho_n\}$ converges to ρ_∞ , namely we have actual convergence and not only just up to conjugacy. Consider the sequence $\{\rho_n \circ \phi_{n*} : \pi_1(M) \rightarrow \text{PSL}(2, \mathbb{C})\}$. By assumption there is a sequence $\{\gamma_n\} \subset \text{PSL}(2, \mathbb{C})$ so that $\{\gamma_n(\rho_n \circ \phi_{n*})\gamma_n^{-1}\}$ converges. Since $\pi_1(N)$ is not Abelian, the non-elementary Kleinian group $\rho_\infty(\pi_1(N))$ contains 2 hyperbolic isometries $\rho_\infty(g_1)$ and $\rho_\infty(g_2)$ with disjoint fixed point sets. By assumption, we have $\rho_n \circ \phi_{n*}(g_i) = \rho_n(g_i)$ for $i = 1, 2$. Thus taking $a_n = \rho_n(g_1)$ and $b_n = \rho_n(g_2)$, we get from Claim 7.1 that, up to extracting a subsequence, $\{\gamma_n\}$ converges. Hence, up to extracting a subsequence $\{\rho_n \circ \phi_{n*}\}$ converges. We have $\rho_n \circ \phi_{n*}(\pi_1(M)) = \rho_n(\pi_1(M))$. Since $\{\rho_n \circ \phi_{n*}\}$ converges and since $\{\rho_n\}$ converges

strongly, for any $a \in \pi_1(M)$ there is $b \in \pi_1(M)$ such that $\{\rho_n \circ \phi_{n*}(a)\}$ converges to $\rho_\infty(b)$. It follows from [Jor] that for n sufficiently large $\phi_{n*}(a) = b$. Thus we have proven that for any closed curve $a \subset M$ there is a closed curve $b \subset M$ such that $\phi_n(a)$ is homotopic to b for n sufficiently large.

Consider a pants decomposition P of $\partial_{\chi < 0}M$. By the above paragraph, for n sufficiently large, ϕ_n maps each leaf c of P to some given curve $d \subset \partial M$ that is freely homotopic to c . By [Joh] and [JaS], for a given curve c , there are only finitely many such simple closed curves d up to isotopy in ∂M . It follows that up to extracting a subsequence there is n_0 such that $\phi_{n_0}^{-1} \circ \phi_n$ maps P to itself up to isotopy. Consider another pants decomposition $P' \subset \partial_{\chi < 0}M$ such that the components of $\partial_{\chi < 0}M - (P \cup P')$ are discs. Using the same arguments we find a diffeomorphism ϕ and a subsequence of $\{\phi_n\}$ such that $\phi^{-1} \circ \phi_n(P)$ is isotopic to P and $\phi^{-1} \circ \phi_n(P')$ is isotopic to P' in ∂M . It follows that the restriction of $\phi^{-1} \circ \phi_n$ to $\partial_{\chi < 0}M$ is isotopic to the identity. By [Joh], up to passing to a subsequence, $\phi^{-1} \circ \phi_n$ is isotopic to the identity for any n . This concludes the proof of Lemma 7.2. $\square$

This is enough to conclude the proof of Theorem C in most cases that remain. Let W be a characteristic submanifold for M . Such a characteristic submanifold is an union of I -bundles and solid tori and its basic property is that any essential annulus in M is isotopic to an annulus lying in W . The existence of such a characteristic submanifold has been proved in [Joh] and [JaS]. If we assume that a component N of $\partial M - \partial W$ has a non-Abelian fundamental group, then we can argue as follows. By [Joh], up to extracting a subsequence there is a diffeomorphism $\phi : M \rightarrow M$ such that each $(\phi \circ \phi_n)|_{M-W}$ is isotopic to the identity. In particular, $(\phi \circ \phi_n)|_N$ is isotopic to the identity. By assumption the sequences $\{\rho_n\}$ and $\{\rho_n \circ (\phi \circ \phi_n)_*\}$ converge in $\text{SH}(M)$. It follows from Lemma 7.2 that up to extracting a subsequence and up to changing ϕ , the diffeomorphism $\phi \circ \phi_n$ is isotopic to the identity for any n . Hence, up to extracting a subsequence, the diffeomorphisms ϕ_n are isotopic and we are done.

For the case where $\partial M - \partial W$ has zero Euler characteristic, we use Theorem D to find an incompressible surface $H \subset \partial M$ satisfying the hypothesis of Lemma 7.2. Let $\Gamma_n = (F_n, m_n, L_n)$ and $\Gamma'_n = (F'_n, m'_n, L'_n)$ be the ending gallimaufries of ρ_n and $\rho'_n = \rho_n \circ \phi_{n*}$. Since we have assumed that ρ_n is minimally parabolic and geometrically finite, we have $F_n = F'_n = \partial_{\chi < 0}M$ and $L_n = L'_n = \emptyset$. Let $\Gamma_\infty = (F_\infty, m_\infty, L_\infty)$ and $\Gamma'_\infty = (F'_\infty, m'_\infty, L'_\infty)$ be the ending gallimaufries of ρ_∞ and ρ'_∞ . By Theorem D, $\{\Gamma_n\}$ converges to Γ_∞ and $\{\Gamma'_n\}$ converges to Γ'_∞ . To each end of $\mathbb{H}^3/\rho_\infty(\pi_1(M))$ corresponds a subsurface of ∂M ; we will say that this subsurface faces an end of $\mathbb{H}^3/\rho_\infty(\pi_1(M))$. Now we reduce the search for H to the search for a curve $c \subset \partial M$ with some specific properties.

Claim 7.3. *Let $c \subset \partial M$ be a simple closed curve such that $\phi_n(c)$ is isotopic to c on ∂M for any n . Consider the incompressible surfaces $H \supset c$ and $H' \supset c$ such that H faces an end of $\mathbb{H}^3/\rho_\infty(\pi_1(M))$ and that H' faces an end of $\mathbb{H}^3/\rho'_\infty(\pi_1(M))$. Assume that c is peripheral neither in H nor in H' . Then there is a diffeomorphism $\phi : M \rightarrow M$ and a subsequence such that each $(\phi^{-1} \circ \phi_n)|_H$ is isotopic to the identity.*

Proof. We show that the ends faced by H and H' are of the same type.

Claim 7.4. *The surface H' faces a geometrically finite end of M'_∞ if and only if H faces a geometrically finite end of M_∞ .*

Proof. By Theorem D, H , respectively H' , faces a geometrically finite end if and only if $\{\ell_{m_n}(c)\}$ is bounded, respectively $\{\ell_{m_n}(\phi_n(c))\}$ is bounded. By assumption, $\phi_n(c)$ is isotopic to c hence we have $\ell_{m_n}(\phi_n(c)) = \ell_{m_n}(c)$. Thus H' faces a geometrically finite end of M'_∞ if and only if H faces a geometrically finite end of M_∞ . $\square$

Consider the case where H and H' face geometrically finite ends. By Theorem D, $\{m_n|_H\}$ and $\{m'_n|_{H'}\}$ converge to complete hyperbolic metrics on H and H' respectively. Since $\phi_n(H)$ contains c , $\phi_n(H)$ intersects H' . But $\ell_{m'_n}(\phi_n(\partial\overline{H})) \rightarrow 0$. Thus, for n large enough, $\phi_n(\partial\overline{H}) \cap \overline{H'} \subset \partial\overline{H'}$ up to isotopy. Similarly, $\phi_n^{-1}(\partial\overline{H'}) \cap \overline{H} \subset \partial\overline{H}$ up to isotopy. This is possible only if $\phi_n(H) = H'$, up to isotopy, for n sufficiently large. Thus up to extracting a subsequence and composing by a fixed diffeomorphism, we may assume $H' = H$ up to isotopy. For a given curve $d \subset H$, $\{\ell_{m'_n}(\phi_n(d)) = \ell_{m_n}(d)\}$ is bounded. Hence there are only finitely many possibilities for its image up to isotopy. It follows that there are only finitely many possibilities for the isotopy class of the diffeomorphism $\phi_n|_H : H \rightarrow H$. This concludes the proof of Claim 7.3 when H faces a geometrically finite end of M_∞ .

Assume now that H faces a geometrically infinite end of M_∞ with ending lamination L and that H' faces a geometrically infinite end of M'_∞ with ending lamination L' . Consider pants decompositions P and P' of $\partial_{\chi < 0} M$ such that $\ell_{m_n}(d) \leq \varepsilon$ for any leaf d of P and $\ell_{m'_n}(d') \leq \varepsilon$ for any leaf d' of P' . Consider $\varepsilon_n \rightarrow 0$ such that, up to extracting a subsequence, $\varepsilon_n \mathcal{F}_P(m_n) \cap H$ tends to a measured geodesic lamination λ . We have $\varepsilon_n \ell_{m'_n}(c) = \varepsilon_n \ell_{m'_n}(\phi_n(c)) = \ell_{m_n}(c) \rightarrow i(\lambda, c)$. Hence, up to extracting a subsequence, $\{\varepsilon_n \mathcal{F}_{P'}(m'_n) \cap H'\}$ tends to a measured geodesic lamination λ' . For a component d of $\partial\overline{H}$, we have $\varepsilon_n \ell_{m_n}(d) \rightarrow 0$. Hence $\varepsilon_n \ell_{m'_n}(\phi_n(d)) \rightarrow 0$. Extract a subsequence such that $\phi_n(d)$ converges in the Hausdorff topology to a geodesic lamination D . Since d is disjoint from c , D is disjoint from c . If D intersects H' without being peripheral, then D would cross $|\lambda'|$. Since $\varepsilon_n \mathcal{F}_{P'}(m'_n) \cap H'$ tends to λ' we would then have $\liminf \varepsilon_n \ell_{m'_n}(\phi_n(d)) = \liminf \varepsilon_n i(\mathcal{F}_{P'}(m'_n), \phi_n(d)) > 0$. Thus we have proved $\phi_n(\partial\overline{H}) \cap \overline{H'} \subset \partial\overline{H'}$ up to isotopy. Similarly we have $\phi_n^{-1}(\partial\overline{H'}) \cap \overline{H} \subset \partial\overline{H}$ up to isotopy and we may assume $H' = H$ up to isotopy.

Let us extend c to a pants decomposition C of H and consider a leaf d of C . Extract a subsequence such that $\phi_n(d)$ converges to a geodesic lamination $D \subset H$. Since d is disjoint from c , D is disjoint from c and crosses $|\lambda|$. If D is not a simple closed curve, we have $\varepsilon_n i(\mathcal{F}_{P'}(m'_n), \phi_n(d)) \rightarrow \infty$. By Theorem 2.2, we have $\varepsilon_n \ell_{m'_n}(\phi_n(d)) \rightarrow \infty$. This would contradict $\varepsilon_n \ell_{m'_n}(\phi_n(d)) = \varepsilon_n \ell_{m_n}(d) \rightarrow i(d, \lambda)$. Thus, up to extracting a subsequence $\phi_n(d)$ does not depend on n . It follows that there is $\phi : M \rightarrow M$ such that up to extracting a subsequence $\phi^{-1} \circ \phi_n$ leaves C invariant, up to isotopy. For each leaf d of C , we choose a transverse t , namely a simple closed curve that crosses d and is disjoint from $C - d$. By the same argument, up to extracting a subsequence $\phi_n(t)$ does not depend on d . It follows that, up to changing ϕ , $(\phi^{-1} \circ \phi_n)|_H$ is isotopic to the identity. $\square$

Now we can conclude the proof of Theorem C by finding a curve c satisfying the assumption of Claim 7.3. Let $W \subset M$ be a characteristic submanifold. By [Joh], up to extracting

a subsequence, there is a diffeomorphism $\phi : M \rightarrow M$ such that $(\phi \circ \phi_n)|_{M-W}$ is isotopic to the identity. As we have seen after the proof of Lemma 7.2, if a component of $M - W$ has a non-Abelian fundamental group, we are done.

Assume that all components of $M - W$ have Abelian fundamental groups. Since M is not an I -bundle, either M is acylindrical and $\text{Mod}(M)$ is finite or $M - W$ is not empty. If $\text{Mod}(M)$ is finite, we are done, so assume that $M - W$ is not empty. The components of $M - W$ are solid tori and I -bundles over tori. Let T_1 be such a component of $M - W$.

If T_1 is an I -bundle over a torus, $\partial T_1 \cap \partial M$ is the union of one torus and some (at least one) annuli or Möbius bands. Since Γ_∞ and Γ'_∞ are acylindrical, any curve c lying in such an annulus or Möbius band lies in the "middle" of an end. Namely for any representation $\rho \in \text{SH}(M)$, c lies in a surface H facing an end of M_ρ and c is not peripheral in $\overline{H}$. Using Lemma 5.2 and Claim 5.5, it is now easy to find pants decomposition P_n such that $\{\ell_{m_n}(P_n)\}$ is bounded and that c satisfies the hypothesis of Lemma 7.3.

If T_1 is a solid torus, $T_1 \cap \partial M$ is the union of annuli. If those annuli are not primitive then we are in the same situation as before. Namely, any curve c lying in such an annulus lies in the "middle" of an end. Hence we can conclude as in the preceding paragraph. If those annuli are primitive, then there are at least 3 of them. Since Γ_∞ and Γ'_∞ are acylindrical at most one simple closed curve $d \subset T_1 \cap \partial M$, respectively $d' \subset T_1 \cap \partial M$, is peripheral in a surface H facing an end of M_∞ , respectively in a surface H' facing an end of M'_∞ . Since $T_1 \cap \partial M$ contains at least 3 non-isotopic simple closed curve, we can chose a simple closed curve $c \neq d$ with $\phi_n(c) \neq d'$ for infinitely many n . Thus c lies in a surface H facing an end of M_∞ and c is not peripheral in $\overline{H}$ and $\phi_n(c)$ lies in a surface H' facing an end of M'_∞ and $\phi_n(c)$ is not peripheral in $\overline{H'}$. It is now easy to find pants decompositions $\{P_n\}$ such that $\{\ell_{m_n}(P_n)\}$ is bounded and that c satisfies the hypothesis of Lemma 7.3.

The surface H produced in Claim 7.3 provides us with an incompressible manifold $N = H$ satisfying the hypothesis of Lemma 7.2. From Lemma 7.2, we conclude that up to extracting a subsequence, the diffeomorphisms ϕ_n are isotopic. This concludes the proof of Theorem C. $\square$

When M is a trivial I -bundle over a closed surface S it is easy to see that Theorem C does not hold. Consider a pseudo-Anosov diffeomorphism $\phi : S \rightarrow S$ with stable lamination λ^+ and unstable lamination λ^- . Consider the diffeomorphism $\phi : M \rightarrow M$ whose projection along the fibers is ϕ . Consider the representation $\rho \in \partial \text{SH}(M)$ which is doubly degenerate and has ending laminations $|\lambda^+|$ on one side and $|\lambda^-|$ on the other side. Then ρ is a fixed point of the action of ϕ on $\text{SH}(M)$.

References

- [Ag] I. Agol, *Tameness of hyperbolic 3-manifolds*, preprint (2004), math.GT/0405568.
- [Ah] L.V. Ahlfors, *Finitely generated Kleinian groups*, Amer. J. Math. 86 (1964), 413-429.
- [AnC] J.W. Anderson, R.D. Canary, *Algebraic limits of Kleinian groups which rearrange the pages of a book*, Invent. Math. 126 (1996), 205-214.
- [ACC] J. W. Anderson, R. D. Canary, D. McCullough, *The topology of deformation spaces of Kleinian groups*, Ann. of Math. (2) 152 (2000), no. 3, 693-741.

- [Be] L. Bers, *An inequality for Riemann surface*, Differential geometry and complex analysis (1985), 87-93.
- [Bon] F. Bonahon, *Bouts des variétés hyperboliques de dimension 3*, Ann. of Math. (2) 124 (1986), 71-158.
- [BoO] F. Bonahon, J.-P. Otal, *Laminations mesurées de plissage des variétés hyperboliques de dimension 3*, Ann. Math. (2) 160 (2005), No.3, 1013-1055.
- [Bow] B. Bowditch, *Some results on the geometry of convex hulls in manifolds of pinched negative curvature*, Comment. Math. Helv. 69 (1994), no. 1, 49-81.
- [Bro] J. F. Brock, *Continuity of Thurston's length function*, Geom. Funct. Anal. 10 (2000), 741-797.
- [BCM] J.F. Brock, R.D. Canary, Y.N. Minsky, *The classification of Kleinian surface groups, II: The Ending Lamination Conjecture*, preprint (2004), math.GT/0412006.
- [Br] K. Bromberg, *The space of Kleinian punctured torus groups is not locally connected*, preprint (2006).
- [BrH] K. Bromberg, J. Holt, *Self-bumping of deformation spaces of hyperbolic 3-manifolds*, J. Differential Geom. 57 (2001), no. 1, 47-65.
- [CaG] D. Calegari, D. Gabai, *Shrinkwrapping and the taming of hyperbolic 3-manifolds*, Jour. Amer. Math. Soc. 19 (2006), no. 2, 385-446.
- [Ca] R.D. Canary, *Ends of hyperbolic 3-manifolds*, J. Amer. Math. Soc. 6 (1993), no. 1, 1-35.
- [CEG] R.D. Canary, D.B.A. Epstein, P. Green, *Notes on notes of Thurston*, Analytical and Geometrical Aspects of hyperbolic Space (1987), 3-92.
- [CaM] R.D. Canary, Y.N. Minsky, *On limits of tame hyperbolic 3-manifolds*, J. Differential Geom. 43 (1996), no. 1, 1-41.
- [EpM] D.B.A. Epstein, A.Marden, *Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces*, Analytical and Geometric Aspects of Hyperbolic Space (1987), 113-253.
- [FLP] A. Fathi, F. Laudenbach, V. Ponearu, *Travaux de Thurston sur les surfaces. Seminaire Orsay, Asterisque No.66-67* (1979).
- [He] D. Hejhal, *Universal covering maps for variable regions*, Math. Z. 137 (1974), 7-20.
- [JaS] W. Jaco, P.B. Shalen, *Seifert fibered spaces in 3-manifolds*, Memoirs Amer. Math. Soc. 220 (1979).
- [Joh] K. Johannson, *Homotopy equivalence of 3-manifolds with boundary*, Lectures Notes in Mathematics 761 (1979).
- [Jor] T. Jørgensen, *On discrete groups of Möbius transformations*, Amer. J. Math. 98 (1976), 739-749.
- [JoM] T. Jørgensen, A. Marden, *Algebraic and geometric convergence of Kleinian groups*, Math. Scand. 66 (1990), no. 1, 47-72.
- [Ka] M. Kapovich, *Hyperbolic manifolds and discrete groups* (2000)
- [Kl] G. Kleinedam, *Strong convergence of kleinian groups*, Geom. Funct. Anal. 15 (2005), no. 2, 416-452
- [KeT] S. Kerckhoff, W. P. Thurston, *Noncontinuity of the action of the modular group at Bers' boundary of Teichmüller space*, Invent. Math. 100 (1990), no. 1, 25-47.
- [Le1] C. Lecuire, *Plissage des variété hyperboliques de dimension 3*, Inventiones Mathematicae 164 (2006), no. 1, 85-141.
- [Le2] C. Lecuire, *Continuity of the bending map*, Ann. Fac. Sci Toulouse Math. , 17 serie 6 (2008), no. 1, 93-119.
- [Lev] G. Levitt, *Foliations and laminations on hyperbolic surfaces*, Topology, 22 (1983), 119-135.
- [Mar] A. Marden, *The geometry of finitely generated Kleinian groups*, Ann. of Math. 99 (1974), 383-462.
- [MaT] K. Matsuzaki, M. Taniguchi, *Hyperbolic Manifolds and Kleinian Groups* (1998).
- [McC] D. McCullough, *Compact submanifolds of 3-manifolds with boundary*, Quart. J. Math. (2) 37 (1986), 299-307.
- [MMS] D. McCullough, A. Miller, G. A. Swarup, *Uniqueness of cores of noncompact 3-manifolds*, J. London Math. Soc. (2) 32 (1985), 548-556.

- [McM] C.T. McMullen, *Hausdorff dimension and conformal dynamics. I. Strong convergence of Kleinian groups*, J. Differential Geom. 51 (1999), no. 3, 471-515.
- [Mi1] Y. N. Minsky, *Kleinian groups and the complex of curves*, Geom. Topol. 4 (2000), 117-148.
- [Mi2] Y.N. Minsky *The classification of Kleinian surface groups, I: Models and bounds*, preprint (2004), math.GT/0302208.
- [MoS1] J.W. Morgan, P.B. Shalen, *Degenerations of Hyperbolic structures I: Valuations, trees and surfaces*, Ann. of Math., 120 (1984), 401-476.
- [MoS2] J.W. Morgan, P.B. Shalen, *Degenerations of Hyperbolic structures III: actions of 3-manifolds groups on trees and Thurston's compactness theorem*, Ann. of Math. 127 (1988), 457-519.
- [Ot1] J.-P. Otal, *Sur la dégénérescence des groupes de Schottky*, Duke Math. J. 74 (1994), 777-792.
- [Ot2] J.-P. Otal, *Le théorème d'hyperbolisation pour les variétés fibrées de dimension 3*, Astérisque 235 (1996).
- [Pe] J.L. Harer, R.C. Penner, *Combinatorics of train tracks*, Annals of Mathematics Studies 125 (1992).
- [Sc] P. Scott, *Compact submanifolds of 3-manifolds*, J. London Math. Soc. (2) 7 (1973), 246-250
- [Sk] R.K. Skora, *Splitting of surfaces*, Bull. Amer. Math. Soc. 23 (1990), 85-90.
- [Su] D. Sullivan, *Quasiconformal homeomorphisms and dynamics. II. Structural stability implies hyperbolicity for Kleinian groups*, Acta Math. 155 (1985), no. 3-4, 243-260.
- [Th1] W.P. Thurston, *The topology and geometry of 3-manifolds*, Lectures Notes, Princeton University, 1976-79.
- [Th2] W.P. Thurston, *Hyperbolic structures on 3-manifolds. I. Deformation of acylindrical manifolds*, Ann. of Math. (2) 124 (1986), no. 2, 203-246.
- [Th3] W.P. Thurston, *Hyperbolic structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle*, preprint arXiv:math.GT/9801045 (1998).
- [Wa] F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, Ann. Math. 87 (1968), 56-88.